

# Construction of two-dimensional quantum field models through Longo-Witten endomorphisms

**Yoh Tanimoto** \*

e-mail: `yoh.tanimoto@theorie.physik.uni-goettingen.de`

Institut für Theoretische Physik, Universität Göttingen

Friedrich-Hund-Platz 1, 37077 Göttingen, Germany.

## Abstract

We present a procedure to construct families of local, massive and interacting Haag-Kastler nets on the two-dimensional spacetime through an operator-algebraic method. An existence proof of local observable is given without relying on modular nuclearity.

By a similar technique, another family of wedge-local nets is constructed using certain endomorphisms of conformal nets recently studied by Longo and Witten.

## 1 Introduction

In a series of papers [18, 33, 6] we have investigated operator-algebraic methods based on conformal field theory to construct quantum field models on two-dimensional spacetime with a weak localization property. Although we succeeded to obtain various examples and general structural results, there was missing one important property: strict localization. In the present paper we construct further new families of quantum field models and prove their strict locality. In other words, we construct interacting two-dimensional Haag-Kastler nets, using techniques from conformal nets.

In recent years the operator-algebraic approach (algebraic QFT) to quantum field theory has seen many developments. A fundamental idea is that the whole quantum field model can be recovered from the set of observables localized in an unbounded wedge-shaped region and the spacetime translations [7]. Conversely, a strategy to construct models is first to obtain models localized in wedges then to prove the existence of fully localized observables. The most successful application of this strategy is the construction of scalar factorizing S-matrix models in two dimensions [22].

---

\*Supported by Deutscher Akademischer Austauschdienst, Hausdorff Institut für Mathematik and Alexander von Humboldt Stiftung, and in part by Courant Research Centre “Higher Order Structures in Mathematics”, Göttingen.

Construction of interacting quantum field theory in four dimensions remains one of the most important open problems in mathematical physics. Further steps toward higher dimensions within the above operator-algebraic approach have been obtained [13, 23], however, the second step to find localized observables has so far turned out to be unsuccessful [13]. Furthermore, we found that conformal covariance implies triviality of scattering in two dimensions [34] yet it is possible to construct weakly localized, conformally covariant models [33]. Thus construction of weakly localized models should be considered as an important but intermediate step.

We have constructed weakly localized massless models using chiral components of two-dimensional conformal field theory [18, 33, 6], however, strict locality is disproved in some of them and open in the others. The only known operator-algebraic technique to find local observables is wedge-splitting [12] (see Section 5). This property is apparently difficult to hold in massless models, especially known to be invalid in conformal case. Actually, one encounters a similar problem in the form factor bootstrap program to integrable models [31]: one starts with a given S-matrix and matrix coefficients of local operators are given as solutions of the so-called form factor equations. One can easily observe that the convergence of these matrix coefficients is much worse in massless case. Hence, in order to avoid such technical difficulty, one should try to construct massive models (besides, the convergence of form factors is a difficult problem even in massive models [1]).

Although our previous constructions are fundamentally based on conformal field theory, one can still find a connection with massive models. We will investigate this issue more systematically in a separate paper [5]. Here we use again Longo-Witten endomorphisms [27] to construct weakly localized massive models. Then we show that some of them are actually strictly local by examining wedge-splitting. Note that in [22] wedge-splitting has been proved through modular nuclearity [12]. We provide both a direct proof of wedge-splitting and a simple proof of modular nuclearity of certain models. Once strict locality is demonstrated, one can apply the standard scattering theory and check that S-matrix is nontrivial. As we will see, this construction can be considered as a generalization of the Federbush model. The Federbush model is an integrable model with S-matrix independent from rapidity [28]. Local observables in this model have been found in [29] for certain range of coupling constant. We construct Haag-Kastler nets which have the same S-matrix as those Federbush models with the coupling constant of arbitrary size. The model has multiple one-particle spectrum, hence the strict locality has not been treated in [22]. We construct another family of wedge-local nets (Borchers triples) using Longo-Witten endomorphisms [27]. Although strict locality is open for this latter family, two constructions look quite similar.

The present construction uses as a starting point the free field or any quantum field which gives a wedge-split net. The new models are constructed on the tensor product of the Hilbert space of the input field (net). One could call our construction a “deformation” of the trivial combination of the input fields. The term “deformation” had a specific sense in [13] and [23], and in massless cases it has been revealed that the connection with free (trivial scattering) models is generic [33]. In addition, one notes that such a “deformation” is always related to a certain structure of the input field. The deformation in [13] exploits

the translation covariance, while the chiral decomposition plays a crucial role in [33, 6]. If one desires further different deformations, it is a natural idea to introduce a further structure to the “undeformed” model. In this paper this idea will be realized using tensor product and inner symmetry. Such structure is easily found in examples. Indeed, our procedure in Section 5 can be applied to the tensor product of Lechner’s models [22] and immediately gives new strict local nets.

This paper is organized as follows: after introductory Sections 1, 2, we first exhibit ways to construct Borchers triples. Section 3 is of general nature: given a Borchers triple with certain symmetry, we show a way to construct new Borchers triples. Section 4 demonstrates how to construct massive Borchers triples out of Longo-Witten endomorphisms of the  $U(1)$ -current net. Then we turn to strict locality in Section 5. We collect arguments to prove strict locality of the models from Section 3. Then we apply them to the concrete cases including the complex massive free field to obtain interacting Haag-Kastler nets (Section 6). The two-particle S-matrix of the models from Section 4 will be also calculated.

## 2 Preliminaries

### 2.1 Two dimensional Haag-Kastler net and Borchers triples

In algebraic QFT, models of quantum field theory are realized as nets of von Neumann algebras [20]. A **Haag-Kastler net (of von Neumann algebras)**  $\mathcal{A}$  on  $\mathbb{R}^2$  is a map  $O \mapsto \mathcal{A}(O)$  from the set of open regions  $\{O\}$  in two dimensional Minkowski space  $\mathbb{R}^2$  to the set of von Neumann algebras on a fixed Hilbert space  $\mathcal{H}$  such that

- (1) **(Isotony)** If  $O_1 \subset O_2$ , then  $\mathcal{A}(O_1) \subset \mathcal{A}(O_2)$ .
- (2) **(Locality)** If  $O_1$  and  $O_2$  are spacelike separated, then  $\mathcal{A}(O_1)$  and  $\mathcal{A}(O_2)$  commute.
- (3) **(Poincaré covariance)** There is a (strongly continuous) unitary representation  $U$  of the (proper orthochronous) Poincaré group  $\mathcal{P}_+^\uparrow$  such that  $U(g)\mathcal{A}(O)U(g)^* = \mathcal{A}(gO)$  for  $g \in \mathcal{P}_+^\uparrow$ .
- (4) **(Positivity of energy)** The joint spectrum of the restriction of  $U$  to the translation subgroup ( $\cong \mathbb{R}^2$ ) is contained in the closed forward lightcone  $V_+ := \{(p_0, p_1) \in \mathbb{R}^2 : p_0 \geq |p_1|\}$ .
- (5) **(Vacuum vector)** There is a (up to scalar) unique vector  $\Omega$  which is invariant under  $U(g)$  and cyclic for  $\mathcal{A}(O)$ .

Note that the condition on the vacuum  $\Omega$  contains the Reeh-Schlieder property, which can be derived if one assumes the additivity of the net. We take this as an axiom for net for simplicity (see the discussion in [35, Section 2]). From these assumptions, the following is automatic [3].

- (6) **(Irreducibility)** The von Neumann algebra  $\bigvee_{O \subset \mathbb{R}^2} \mathcal{A}(O)$  is equal to  $B(\mathcal{H})$ .

Precisely, the triple  $(\mathcal{A}, U, \Omega)$  should be called a Haag-Kastler net, however we say  $\mathcal{A}$  is a net if no confusion arises. If one has a net  $\mathcal{A}$ , under certain conditions one can define S-matrix (see Section 6). The main objective in this paper is to construct examples of two-dimensional Haag-Kastler nets with nontrivial S-matrix.

Yet, it appears very difficult to construct such nets from a scratch. Fortunately, Borchers showed that it suffices to have a triple of a *single* von Neumann algebra associated with the wedge-shaped region  $W_R := \{a \in \mathbb{R}^2 : a_1 > |a_0|\}$ , the spacetime translations and the vacuum. More precisely, a **Borchers triple** is a triple  $(\mathcal{M}, T, \Omega)$  of a von Neumann algebra  $\mathcal{M}$ , a unitary representation  $U$  of  $\mathbb{R}^2$  and a vector  $\Omega$  such that

- (1) If  $a \in W_R$ , then  $\text{Ad } T(a)(\mathcal{M}) \subset \mathcal{M}$ .
- (2) The joint spectrum of  $T$  is contained in  $V_+$ .
- (3)  $\Omega$  is cyclic and separating for  $\mathcal{M}$ .

The correspondence between nets and Borchers triples is as follows (see [7, 22]): If  $(\mathcal{A}, U, \Omega)$  is a (two-dimensional) Haag-Kastler net, then  $(\mathcal{A}(W_R), U|_{\mathbb{R}^2}, \Omega)$  is a Borchers triple, where  $U$  is restricted to the translation subgroup  $\cong \mathbb{R}^2$ .

On the other hand, if  $(\mathcal{M}, T, \Omega)$  is a Borchers triple, then one can define a net as follows: in the two-dimensional Minkowski space any double cone  $D$  is represented as the intersection of two wedges  $D = (W_R + a) \cap (W_L + b)$ . Then we define for double cones  $\mathcal{A}(D) := \text{Ad } T(a)(\mathcal{M}) \cap \text{Ad } T(b)(\mathcal{M}')$ . For a general region  $O$  we take  $\mathcal{A}(O) := \bigvee_{D \subset O} \mathcal{A}(D)$ , where the union runs all the double cones contained in  $O$ . Then one can observe that  $\mathcal{A}$  satisfies isotony and locality. Borchers further proved that the representation  $T$  of  $\mathbb{R}^2$  extends to a representation  $U$  of  $\mathcal{P}_+^\uparrow$  through Tomita-Takesaki theory [7] such that  $\mathcal{A}$  is covariant and  $\Omega$  is invariant under  $U$  (the representation of boosts is given by the modular group). Positivity of energy is inherited from  $T$  of the Borchers triple. The only missing property is cyclicity of  $\Omega$  for local algebras  $\mathcal{A}(O)$ . Indeed, there are examples of Borchers triples which fail to satisfy this cyclicity [33, Theorem 4.16].

Hence, a general strategy to construct Haag-Kastler nets is first to construct Borchers triples and then to check cyclicity of the vacuum. This program has been first completed in [22]. Note that the latter condition is actually very hard to check directly. In [22], Lechner proved instead the so-called modular nuclearity [12], which is a sufficient condition for the cyclicity of the vacuum. In this paper, we will provide new examples of Borchers triples for which the cyclicity of the vacuum can be proved without relying on modular nuclearity (see Section 5.2). We say that a Borchers triple is **strictly local** if  $\Omega$  is cyclic for  $\mathcal{A}(O)$  constructed as above. A strictly local Borchers triple corresponds to a Haag-Kastler net.

If there is a unitary operator  $V$  which commutes with  $U(g)$  and if  $\text{Ad } V$  preserves each local algebra  $\mathcal{A}(O)$ , then we call  $\text{Ad } V$  an **inner symmetry** of the net  $\mathcal{A}$ . By the uniqueness of  $\Omega$ ,  $\text{Ad } V$  preserves the vacuum state  $\langle \Omega, \cdot \Omega \rangle$ . By the irreducibility, the automorphism  $\text{Ad } V$  is implemented uniquely by  $V$  up to a scalar. We require always that  $V\Omega = \Omega$ . By this requirement, the implementation is unique. A general strategy to use them in order to construct new Borchers triples will be explained in Section 3 and concrete examples of

inner symmetry will be discussed in Section 6.1. A collection of inner symmetries may form a group  $G$ . In such a case we say that  $G$  acts on the net by inner symmetry. Similarly one can consider an action of  $V$  on Borchers triples. In this case, one says that  $V$  implements an inner symmetry if  $\text{Ad } V$  preserves  $\mathcal{M}$  and commutes with  $T$ . We say also that  $G$  acts by inner symmetry when such  $V$ 's form a group.

## 2.2 Conformal nets and Longo-Witten endomorphisms

In some of our constructions, the main ingredients come from chiral conformal field theory. Let us summarize here its operator-algebraic treatment (see also [19, 26]). An open, connected, non dense and non empty subset  $I$  of  $S^1$  is called an **interval** in  $S^1$ . We identify  $\mathbb{R}$  as a dense subset in  $S^1$  by the stereographic projection. The Möbius group  $\text{PSL}(2, \mathbb{R})$  acts on  $S^1 = \mathbb{R} \cup \{\infty\}$  and under this identification it contains translations and dilations of  $\mathbb{R}$ . A **(Möbius covariant) net (of von Neumann algebras) on  $S^1$**  is an assignment of von Neumann algebras  $\mathcal{A}_0(I)$  on a common Hilbert space  $\mathcal{H}_0$  to intervals  $I$  such that

- (1) **(Isotony)** If  $I_1 \subset I_2$ , then  $\mathcal{A}_0(I_1) \subset \mathcal{A}_0(I_2)$ .
- (2) **(Locality)** If  $I_1$  and  $I_2$  are disjoint, then  $\mathcal{A}_0(I_1)$  and  $\mathcal{A}_0(I_2)$  commute.
- (3) **(Möbius covariance)** There is a (strongly continuous) unitary representation  $U_0$  of  $\text{PSL}(2, \mathbb{R})$  such that  $U_0(g)\mathcal{A}_0(I)U_0(g)^* = \mathcal{A}_0(gI)$  for  $g \in \text{PSL}(2, \mathbb{R})$ .
- (4) **(Positivity of energy)** The generator of the subgroup of translation in the representation  $U_0$  has positive spectrum.
- (5) **(Vacuum vector)** There is a (up to scalar) unique vector  $\Omega_0$  which is invariant under  $U_0(g)$  and cyclic for  $\mathcal{A}_0(I)$ .

As in two dimensions, we call  $\mathcal{A}_0$  a net, however the actual object of interest is the triple  $(\mathcal{A}_0, U_0, \Omega_0)$ . From these assumptions many properties automatically follow, among which of importance in our application are

- (6) **(Irreducibility)** The von Neumann algebra  $\bigvee_{I \subset S^1} \mathcal{A}_0(I)$  is equal to  $B(\mathcal{H}_0)$ .
- (7) **(Haag duality on  $S^1$ )** It holds that  $\mathcal{A}_0(I)' = \mathcal{A}_0(I')$ , where  $I'$  denotes the interior of the complement of  $I$  in  $S^1$ .
- (8) **(Bisognano-Wichmann property)** The modular group  $\Delta_0^{it}$  of  $\mathcal{A}_0(\mathbb{R}_+)$  associated with  $\Omega_0$  is equal to  $U_0(\delta(-2\pi t))$ , where  $\delta$  is the dilation in  $\text{PSL}(2, \mathbb{R})$ .

An **inner symmetry** of a conformal net  $\mathcal{A}_0$  is a collection of automorphisms of each local algebra  $\mathcal{A}_0(O)$  implemented by a common unitary operator  $V_0$  which preserves the vacuum state  $\langle \Omega_0, \cdot \Omega_0 \rangle$ . In conformal (Möbius covariant) case, it automatically follows that  $V_0$  commutes with  $U_0(g)$  thanks to Bisognano-Wichmann property.

A **Longo-Witten endomorphism** of  $\mathcal{A}_0$  is an endomorphism of  $\mathcal{A}_0(\mathbb{R}_+)$ , implemented by a unitary operator  $V_0$  which commutes with  $U_0(g)$ , where  $g$  is a translation. An inner symmetry restricted to  $\mathcal{A}_0(\mathbb{R}_+)$  is a Longo-Witten endomorphism.

We will give a concrete example of conformal net and of Longo-Witten endomorphisms in Section 4.

## 2.3 The massive scalar free field

Our main construction strategy is based on simpler examples with certain properties. Let us quickly review the simplest quantum field.

The free field is constructed from an irreducible representation of the Poincaré group through second quantization. We use the notation which is to some extent consistent with [25]. The one-particle Hilbert space is  $\mathcal{H}_1 := L^2(\mathbb{R}, d\theta)$  and for the mass  $m > 0$  the (proper orthochronous) Poincaré group acts by  $U_1(a, \lambda)\psi(\theta) = e^{ip(\theta) \cdot a}\psi(\theta - \lambda)$ , where  $p(\theta) := (m \cosh(\theta), m \sinh(\theta))$  parametrizes the mass shell. We introduce the (auxiliary) unsymmetrized Hilbert space  $\mathcal{H}^\Sigma := \bigoplus \mathcal{H}_1^{\otimes n}$  and the (physical) symmetrized Hilbert space  $\mathcal{H}_r := \bigoplus Q_n \mathcal{H}_1^{\otimes n}$ , where  $Q_n$  is the projection onto the symmetric subspace.

Let us denote the  $n$ -th component of a vector  $\Psi \in \mathcal{H}_r$  by  $(\Psi)_n$ . For a vector  $\psi \in \mathcal{H}_1$  and  $\Psi$  which has only finitely many components, the creation operator  $a^\dagger$  is defined by  $(a^\dagger(\psi)\Psi)_n = Q_n(\psi \otimes \Psi_{n-1})$ . The annihilation operator is the adjoint  $a(\psi) = a^\dagger(\psi)^*$ . With this notation,  $a^\dagger$  is linear and  $a$  is antilinear with respect to  $\psi$ . The free quantum field  $\phi$  is now given by

$$\phi(f) := a^\dagger(f^+) + a(J_1 f^-), \quad f^\pm(\theta) = \frac{1}{2\pi} \int d^2 a f(a) e^{\pm i p(\theta) \cdot a},$$

where  $f$  is a test function  $\mathcal{S}(\mathbb{R}^2)$  and  $J_1 \psi(\theta) = \overline{\psi(\theta)}$ . This field is local, in the sense that if  $f$  and  $g$  have spacelike separated supports, then  $\phi(f)$  and  $\phi(g)$  commute on an appropriate domain.

Finally we introduce the free field net. For each open region  $O \subset \mathbb{R}^2$ , we set

$$\mathcal{A}_r(O) := \{e^{i\phi(f)} : \text{supp } f \subset O\}''.$$

We have the second quantized representation  $U_r := \Gamma(U_1)$  and the Fock vacuum vector  $\Omega_r \in \mathcal{H}_r$ . This triple  $(\mathcal{A}_r, U_r, \Omega_r)$  is a two-dimensional Haag-Kastler net and referred to as the **free massive scalar net**. The subscript  $r$  is intended for the *real* scalar field. Furthermore, this net satisfies the modular nuclearity [12, Section 4], a property which we will explain in more detail in Section 5.

## 2.4 Scalar factorizing S-matrix models

The free net has many important features, but is not interacting. Lechner has constructed a large family of interacting nets [22]. Here we only briefly summarize the construction and their fundamental properties.

Let  $S_2(\theta)$  be a bounded analytic function on the strip  $0 < \Im \theta < \pi$ , continuous on the boundary, with the property

$$S_2(\theta)^{-1} = \overline{S_2(\theta)} = S_2(-\theta) = S_2(\theta + i\pi)$$

for  $\theta \in \mathbb{R}$ . This is called the two-particle scattering function.

This time again the construction of the (wedge-local) field and the net is based on the one-particle space  $\mathcal{H}_1$  above. For  $\Psi_n \in \mathcal{H}_1^{\otimes n}$ , the  $S_2$ -permutation is given by

$$(D_{S_2,n}(\tau_j)\Psi)(\theta_1, \dots, \theta_n) = S_2(\theta_{j+1} - \theta_j)\Psi(\theta_1, \dots, \theta_{j+1}, \theta_j, \dots, \theta_n).$$

Let  $Q_{S_2,n}$  be the orthogonal projection onto the subspace of  $\mathcal{H}_1^{\otimes n}$  invariant under  $\{D_{S_2,n}(\tau_j) : 1 \leq j \leq n-1\}$ . The full Hilbert space is  $\mathcal{H}_{S_2} := \bigoplus Q_{S_2,n} \mathcal{H}_1^{\otimes n}$  and the representation  $U_1$  promotes to  $U_{S_2}$  on  $\mathcal{H}_{S_2}$  by the second quantization. We define similarly the creation and annihilation operators  $(z_{S_2}^\dagger(\psi)\Phi)_n = Q_{S_2,n}(\psi \otimes \Phi_{n-1})$  and  $z_{S_2}(\psi) = z_{S_2}^\dagger(\psi)^*$ . The quantum field is defined accordingly by

$$\phi_{S_2}(f) := z_{S_2}^\dagger(f^+) + z_{S_2}(J_1 f^-), \quad f^\pm = \frac{1}{2\pi} \int d^2 a f(a) e^{\pm i p(\theta) \cdot a},$$

but this time  $\phi_{S_2}$  is only wedge-local, hence the net is defined through Borchers triple.

The von Neumann algebra  $\mathcal{M}_{S_2}$  is defined by

$$\mathcal{M}_{S_2} := \{e^{i\phi_{S_2}(f)} : \text{supp } f \subset W_L\}'$$

(note that here we take only the single commutant and  $f$  has support in  $W_L$ , while  $\mathcal{M}_{S_2}$  corresponds to  $W_R$ . By the wedge-duality, this is just a matter of convention (see [25])). The triple  $(\mathcal{M}_{S_2}, U_{S_2}, \Omega_{S_2})$  is a Borchers triple [21]. Furthermore, the modular nuclearity holds if  $S_2$  is regular and fermionic [22], hence the triple is wedge-split in those cases.

### 3 Borchers triples through inner symmetries

In this Section we make the first step in our main construction. We start with a Borchers triple with inner symmetry and construct new triples. Note that, differently from our previous results [33, 6], the new triples are defined on a different Hilbert space although the formulae look very similar.

First we state a key lemma, which can be obtained as a special case of [33, Lemma 4.1].

**Lemma 3.1.** *Let  $\mathcal{M}_c$  be a von Neumann algebra and  $Q_c$  a self-adjoint operator such that  $\text{sp } Q_c \subset \mathbb{Z}$  and  $\text{Ad } e^{i2\pi\kappa Q_c}(\mathcal{M}_c) = \mathcal{M}_c$  for any  $\kappa \in \mathbb{R}$ . Then, for  $\kappa \in \mathbb{R}$ ,  $x \otimes \mathbb{1}$  commutes with  $\text{Ad } e^{i2\pi\kappa Q_c \otimes Q_c}(x' \otimes \mathbb{1})$  where  $x \in \mathcal{M}_c$  and  $x' \in \mathcal{M}_c'$ .*

*Proof.* For any  $\kappa \in \mathbb{R}$ ,  $x$  commutes with  $\text{Ad } e^{i2\pi\kappa Q_c}(x')$  by assumption, hence the commutativity lemma [33, Lemma 4.1] applies.  $\square$

Analogously one can prove the following.

**Lemma 3.2.** *Let  $N$  be an integer,  $\mathcal{M}_c$  a von Neumann algebra and  $Q_c$  a self-adjoint operator such that  $\text{sp } Q_c \subset \mathbb{Z}$  and  $\text{Ad } e^{i\frac{2\pi k}{N}Q_c}(\mathcal{M}_c) = \mathcal{M}_c$  for any  $k \in \mathbb{Z}$ . Then, for  $k \in \mathbb{Z}$ ,  $x \otimes \mathbb{1}$  commutes with  $\text{Ad } e^{i\frac{2\pi k}{N}Q_c \otimes Q_c}(x' \otimes \mathbb{1})$  where  $x \in \mathcal{M}_c$  and  $x' \in \mathcal{M}'_c$ .*

Now let  $(\mathcal{M}_c, T_c, \Omega_c)$  be a Borchers triple with an action of  $S^1$  or the finite cyclic group  $\mathbb{Z}_N$  by inner symmetry. The argument of this Section goes parallelly for both groups. There is a unique unitary operator  $V_c(\kappa)$  which implements the inner symmetry by  $S^1$ . For a  $\mathbb{Z}_N$ -action we take  $V_c$  such that  $V_c^k$  implements the inner symmetry for  $k \in \mathbb{Z}_N$ . One can choose  $Q_c$  such that  $V_c(\kappa) = e^{i2\pi\kappa Q_c}$  or  $V_c = e^{i\frac{2\pi}{N}Q_c}$ , respectively. For an action of  $S^1$ ,  $Q_c$  is unique. For  $\mathbb{Z}_N$ , we remark that we can concretely choose  $Q_c$  as follows: for each integer  $j \in [0, N-1]$  we put  $\hat{V}_c(j) := \frac{1}{N} \sum_k e^{-i\frac{2\pi jk}{N}} V_c(k)$ . It is clear that  $\hat{V}_c(j)$  is an orthogonal projection and  $\hat{V}_c(j)\hat{V}_c(l) = 0$  if  $j \not\equiv l \pmod{N}$ . In other words,  $k \mapsto V_c^k$  is a representation of  $\mathbb{Z}_N$ , whose dual group is also  $\mathbb{Z}_N$  and we extract the spectral projections. Then we can take  $Q_c := \sum_j j \hat{V}_c(j)$ . By this construction, it is clear that  $Q_c$  commutes with the translation  $T_c$ . An operator  $Q_c$  with the properties specified above is not unique since one can amplify each spectral component by  $N$ , but if we consider the operator  $e^{i\frac{2\pi k}{N}Q_c \otimes Q_c}$ , it does not depend on the choice of  $Q_c$  (see also the remark after [33, Theorem 4.17]). One sees that  $\text{sp } Q_c \subset \mathbb{Z}$  since  $\mathbb{1} = V_c(1) = e^{i2\pi Q_c}$  for an  $S^1$ -action or  $\mathbb{1} = V_c^N = e^{i2\pi Q_c}$  for a  $\mathbb{Z}_N$ -action (or directly by our choice).

Now we turn to the construction of Borchers triples. Our objects act on the tensor product Hilbert space  $\tilde{\mathcal{H}}_c := \mathcal{H}_c \otimes \mathcal{H}_c$ . Let us denote  $\tilde{V}_{c,\kappa} = e^{i2\pi\kappa Q_c \otimes Q_c}$  for an  $S^1$ -action and  $\tilde{V}_{c,k} = e^{i\frac{2\pi k}{N}Q_c \otimes Q_c}$  for a  $\mathbb{Z}_N$ -action. The tensor product representation  $\tilde{T}_c(a) := T_c(a) \otimes T_c(a)$  has positive spectrum and preserves the new vacuum  $\tilde{\Omega}_c := \Omega_c \otimes \Omega_c$ . The von Neumann algebra is given respectively by

$$\begin{aligned}\tilde{\mathcal{M}}_{c,\kappa} &:= \{x \otimes \mathbb{1}, \text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes y) : x, y \in \mathcal{M}_c\}'' , \\ \tilde{\mathcal{M}}_{c,k} &:= \{x \otimes \mathbb{1}, \text{Ad } \tilde{V}_{c,k}(\mathbb{1} \otimes y) : x, y \in \mathcal{M}_c\}'' .\end{aligned}$$

Note the difference from our previous construction [33]: first of all, this time the input is the two-dimensional Borchers triple, while we used chiral components in [33]. Accordingly, the representation  $\tilde{T}_c$  is now just the two copies of the given one. The formula for  $\tilde{\mathcal{M}}_{c,\kappa}$  or  $\tilde{\mathcal{M}}_{c,k}$  is also similar, but this time we take both  $x$  and  $y$  from the same algebra  $\mathcal{M}_c$ .

**Theorem 3.3.** *The triples  $(\tilde{\mathcal{M}}_{c,\kappa}, \tilde{T}_c, \tilde{\Omega}_c)$  and  $(\tilde{\mathcal{M}}_{c,k}, \tilde{T}_c, \tilde{\Omega}_c)$  are Borchers triples for  $\kappa \in \mathbb{R}$  and  $k \in \mathbb{Z}_N$ , respectively.*

*Proof.* We give a proof only for  $(\tilde{\mathcal{M}}_{c,\kappa}, \tilde{T}_c, \tilde{\Omega}_c)$  constructed from an  $S^1$ -action. The other case is completely parallel.

As in [33, Theorem 4.17], the properties of  $\tilde{T}_c$  and  $\tilde{\Omega}_c$  are readily checked. As for the relation  $\text{Ad } \tilde{T}_c(a)(\tilde{\mathcal{M}}_{c,\kappa}) \subset \tilde{\mathcal{M}}_{c,\kappa}$  for  $a \in W_R$ , one can prove this by noting that  $\tilde{V}_{c,\kappa}$  and  $\tilde{T}_c(a)$  commute and the assumptions that  $\mathcal{M}_c$  and  $T_c$  have the relation.

The cyclicity of  $\tilde{\Omega}_c$  for  $\tilde{\mathcal{M}}_{c,\kappa}$  follows from the cyclicity of  $\Omega_c$  for  $\mathcal{M}_c$  and the fact that  $\tilde{V}_{c,\kappa}(\mathbb{1} \otimes y)\tilde{\Omega}_c = (\mathbb{1} \otimes y)\tilde{\Omega}_c$ . To show the separating property, we need again to prepare a

sufficiently big algebra in the commutant:

$$\tilde{\mathcal{M}}_{c,\kappa}^1 := \{\text{Ad } \tilde{V}_{c,k}(x' \otimes \mathbb{1}), \mathbb{1} \otimes y' : x', y' \in \mathcal{M}'_c\}''.$$

It is easy to see that  $\tilde{\Omega}_c$  is cyclic for  $\tilde{\mathcal{M}}_{c,\kappa}^1$ . In order to see that  $\tilde{\Omega}_c$  is separating for  $\tilde{\mathcal{M}}_{c,\kappa}$ , it is enough to prove that  $\tilde{\mathcal{M}}_{c,\kappa}$  and  $\tilde{\mathcal{M}}_{c,\kappa}^1$  commute. This follows from Lemma 3.1, as in [33, Theorem 4.2].  $\square$

At the end of this Section, we remark that the existence of inner symmetry is not at all exceptional. Indeed, if  $(\mathcal{M}_c, T_c, \Omega_c)$  is a (wedge-split) Borchers triple, then the tensor product  $(\mathcal{M}_c \otimes \mathcal{M}_c, T_c \otimes T_c, \Omega_c \otimes \Omega_c)$  has the flip automorphism which commutes with  $T_c \otimes T_c$  and preserves  $\Omega_c \otimes \Omega_c$ , hence it is an inner symmetry  $\mathbb{Z}_2$ . We present more examples in Section 6.1.

## 4 Borchers triples through Longo-Witten endomorphisms on the $U(1)$ -current net

Here we exhibit another procedure to produce Borchers triples in a more concrete way. We take the free massive net as the starting point (Section 2.3). The formulae are quite similar to those in [33], but should not be confused. Strict locality of the models constructed here is not investigated in the present paper.

### 4.1 Reduction to lightray

Let  $(\mathcal{A}_r, U_r, \Omega_r)$  be the free massive net. The representation  $U_r$  can be restricted to the positive lightray, which we denote by  $U_r^+$  and we obtain a one-dimensional Borchers triple  $(\mathcal{M}_r, U_r^+, \Omega_r)$ , namely a von Neumann algebra  $\mathcal{M}_r := \mathcal{A}_r(W_R)$ , a positive energy representation  $U_r^+$  of  $\mathbb{R}$  and a cyclic separating vector  $\Omega_r$  for  $\mathcal{A}_r$  invariant under  $U_r^+$  such that  $\text{Ad } U_r^+(t)(\mathcal{M}_r) \subset \mathcal{M}_r$  for  $t \in \mathbb{R}_+$ . We denote the negative lightlike translation by  $U_r^-$ .

Let us recall the  $U(1)$ -current net  $(\mathcal{A}^{(0)}, U_0, \Omega_0)$ , which is a conformal net. For its definition, see our previous discussion [33, Section 5]. The point is that the Hilbert space is naturally isomorphic to the Fock space  $\mathcal{H}_r$  of the massive free net, whose one-particle space is  $L^2(\mathbb{R}, d\theta)$  and one considers the second quantization operators. The above one-dimensional triple  $(\mathcal{M}_r, U_r^+, \Omega_r)$  is actually unitarily equivalent to the triple  $(\mathcal{A}^{(0)}(\mathbb{R}_+), T_0, \Omega_0)$ , where  $(\mathcal{A}^{(0)}, U_0, \Omega_0)$  is the  $U(1)$ -current net and  $T_0$  is the restriction of  $U_0$  to the translation subgroup. This will be explained in more detail in [5].

In particular, we can exploit the Longo-Witten endomorphisms found in [27]. Recall that, for an inner symmetric function  $\varphi(z)$ , namely the boundary value on  $\mathbb{R}$  of a bounded analytic function on  $0 < \Im z < \pi$ , one considers the operator  $V_\varphi := \Gamma(\varphi(P_1))$ , where  $P_1$  is the generator of the restriction of  $U_r^+$  on the one-particle space  $\mathcal{H}_1$ ,  $\varphi(P_1)$  denotes a functional calculus and  $\Gamma$  is the second quantization (note that in general the order of second quantization and functional calculus cannot be exchanged:  $\Gamma(\varphi(P_1)) \neq \varphi(\Gamma(P_1))$ ).

Then  $\text{Ad } V_\varphi$  preserves  $\mathcal{M}_r$  and  $V_\varphi$  commutes with  $U_r^+$ . Furthermore,  $V_\varphi$  commutes with  $U_r^-$  since  $U_r^-(a) = \Gamma\left(\exp\left(\frac{it}{P_1}\right)\right)$ , as we see in [5].

## 4.2 Construction of Borchers triples

We work on the tensor product Hilbert space  $\tilde{\mathcal{H}}_r := \mathcal{H}_r \otimes \mathcal{H}_r$ . We fix an inner symmetric function  $\varphi$ . As above,  $P_1$  is the one-particle lightlike translation. Let us recall our argument [33, Section 5].

The physical Hilbert space  $\mathcal{H}_r$  is included in the unsymmetrized Fock space  $\mathcal{H}^\Sigma$ . We consider  $m$  commuting operators on  $\mathcal{H}_1^{\otimes m}$ :

$$\{\mathbb{1} \otimes \cdots \otimes \underset{j\text{-th}}{P_1} \otimes \cdots \otimes \mathbb{1} : 1 \leq j \leq m\}.$$

For  $1 \leq j \leq m$  and  $1 \leq k \leq n$ , let us define operators on  $\mathcal{H}^m \otimes \mathcal{H}^n$ :

$$P_{j,k}^{m,n} := \left( \mathbb{1} \otimes \cdots \otimes \underset{j\text{-th}}{\frac{1}{P_1}} \otimes \cdots \otimes \mathbb{1} \right) \otimes \left( \mathbb{1} \otimes \cdots \otimes \underset{k\text{-th}}{P_1} \otimes \cdots \otimes \mathbb{1} \right)$$

$$R_\varphi^{m,n} := \prod_{j,k} \varphi(P_{j,k}^{m,n}),$$

where  $\varphi(P_{j,k}^{m,n})$  is the functional calculus on  $\mathcal{H}_1^{\otimes m} \otimes \mathcal{H}_1^{\otimes n}$ . Now, our key operator on the unsymmetrized space  $\mathcal{H}^\Sigma \otimes \mathcal{H}^\Sigma$  is

$$\tilde{R}_\varphi := \bigoplus_{m,n} R_\varphi^{m,n} = \bigoplus_{m,n} \prod_{j,k} \varphi(P_{j,k}^{m,n}),$$

where for  $m = 0$  or  $n = 0$  we set  $R_{j,k}^{m,n} = \mathbb{1}$  as a convention. It is easy to see that  $\tilde{R}_\varphi$  naturally restricts to partially symmetrized subspaces  $\mathcal{H}_r \otimes \mathcal{H}^\Sigma$  and  $\mathcal{H}^\Sigma \otimes \mathcal{H}_r$  and to the totally symmetrized space  $\mathcal{H}_r \otimes \mathcal{H}_r$ .

Let  $E_1 \otimes E_1 \otimes \cdots \otimes E_1$  be the joint spectral measure of  $\{\mathbb{1} \otimes \cdots \otimes \underset{k\text{-th}}{P_1} \otimes \cdots \otimes \mathbb{1} : 1 \leq k \leq n\}$ .

One obtains the following expression:

$$\begin{aligned} \varphi(P_{j,k}^{m,n}) &= \int \left( \mathbb{1} \otimes \cdots \otimes \underset{j\text{-th}}{\varphi\left(\frac{p_k}{P_1}\right)} \otimes \cdots \otimes \mathbb{1} \right) \otimes \left( \mathbb{1} \otimes \cdots \otimes \underset{k\text{-th}}{dE_1(p_k)} \otimes \cdots \otimes \mathbb{1} \right) \\ &= \int \left( \mathbb{1} \otimes \cdots \otimes \underset{j\text{-th}}{dE_1(p_j)} \otimes \cdots \otimes \mathbb{1} \right) \otimes \left( \mathbb{1} \otimes \cdots \otimes \underset{k\text{-th}}{\varphi\left(\frac{P_1}{p_j}\right)} \otimes \cdots \otimes \mathbb{1} \right) \end{aligned}$$

Similarly to our previous case [33, Section 5.2], we decompose  $\tilde{R}_\varphi$  with respect only to the right or left component:

$$\begin{aligned}\tilde{R}_\varphi &= \bigoplus_n \int \prod_k \Gamma \left( \varphi \left( \frac{p_k}{P_1} \right) \right) \otimes dE_1(p_1) \otimes \cdots \otimes dE_1(p_n) \\ &= \bigoplus_m \int dE_1(p_1) \otimes \cdots \otimes dE_1(p_m) \otimes \prod_j \Gamma \left( \varphi \left( \frac{P_1}{p_j} \right) \right)\end{aligned}$$

For the proof, we refer to [33, Section 5.2]. The expression in the first line naturally restricts to the partially symmetrized space  $\mathcal{H}_r \otimes \mathcal{H}^\Sigma$  and the second expression to  $\mathcal{H}^\Sigma \otimes \mathcal{H}_r$ .

We have a variant of [33, Lemma 5.2].

**Lemma 4.1.** *It holds for  $x \in \mathcal{A}_r(W_R)$  and  $x' \in \mathcal{A}_r(W_R)'$  that*

$$[x \otimes \mathbb{1}, \text{Ad } \tilde{R}_\varphi(x' \otimes \mathbb{1})] = 0$$

*on the Hilbert space  $\mathcal{H}_r \otimes \mathcal{H}_r$ . Similarly, for  $y \in \mathcal{A}_r(W_R)$  and  $y' \in \mathcal{A}_r(W_R)'$  one has*

$$[\text{Ad } \tilde{R}_\varphi(\mathbb{1} \otimes y), \mathbb{1} \otimes y'] = 0.$$

*Proof.* The operator  $\tilde{R}_\varphi$  is disintegrated into second quantization operators as we saw above. First we consider the first of the commutators above. The operator  $\tilde{R}_\varphi$  restricts naturally to  $\mathcal{H}_r \otimes \mathcal{H}^\Sigma$  and  $x \otimes \mathbb{1}$  and  $x' \otimes \mathbb{1}$  extend naturally to  $\mathcal{H}_r \otimes \mathcal{H}^\Sigma$ . The statement is equivalent to

$$[\text{Ad } \tilde{R}_{\overline{\varphi}}(x \otimes \mathbb{1}), x' \otimes \mathbb{1}] = 0.$$

It is easy to see that if  $\varphi(z)$  is an inner symmetric function, then so are  $\overline{\varphi(1/\overline{z})}$  and hence  $\overline{\varphi(p_k/\overline{z})}$  for  $p_k \geq 0$ . Let us calculate the commutation relation on  $\mathcal{H}_r \otimes \mathcal{H}^\Sigma$ .

$$\text{Ad } \tilde{R}_{\overline{\varphi}}(x \otimes \mathbb{1}) = \bigoplus_n \int \text{Ad} \left( \prod_k \Gamma \left( \overline{\varphi} \left( \frac{p_k}{P_1} \right) \right) \right) (x) \otimes dE_1(p_1) \otimes \cdots \otimes dE_1(p_n).$$

And this commutes with  $x \otimes \mathbb{1}$ . Indeed, since  $x \in \mathcal{A}_r(W_R) = \mathcal{A}_0(\mathbb{R}_+)$  and  $x' \in \mathcal{A}_r(W_R)' = \mathcal{A}_0(\mathbb{R}_-)$ , it follows that  $\text{Ad } \Gamma \left( \overline{\varphi} \left( \frac{p_k}{P_1} \right) \right) (x) \in \mathcal{A}_0(\mathbb{R}_+)$  for any  $p_k \geq 0$  by the result of Longo and Witten [27] (see also the beginning of this Section), and by the fact that the spectral support of  $E_1$  is positive. Then the equality naturally restricts to  $\mathcal{H}_r \otimes \mathcal{H}_r$ .

The proof of the second commutator goes more simply. We only have to consider the expression

$$\text{Ad } \tilde{R}_\varphi(\mathbb{1} \otimes y) = \bigoplus_m \int dE_1(p_1) \otimes \cdots \otimes dE_1(p_m) \otimes \text{Ad} \left( \prod_j \Gamma \left( \varphi \left( \frac{P_1}{p_j} \right) \right) \right) (y).$$

The rest of the argument is parallel as above. □

Let  $T_r$  be the restriction of  $U_r$  to translation (not to be confused with trace). Our Borchers triple is given as follows.

**Theorem 4.2.** *The triple*

- $\tilde{\mathcal{M}}_{r,\varphi} := \{x \otimes \mathbb{1}, \text{Ad } \tilde{R}_\varphi(\mathbb{1} \otimes y) : x, y \in \mathcal{A}_r(W_R)\}''$
- $\tilde{T}_r = T_r \otimes T_r$
- $\tilde{\Omega}_r = \Omega_r \otimes \Omega_r$

*is a Borchers triple.*

*Proof.* The conditions on  $\tilde{T}_r$  and  $\tilde{\Omega}_r$  are readily satisfied since they are same as the tensor product net. The operators  $\tilde{R}_\varphi$  and  $\tilde{T}_r$  commute since both are the functional calculus of the same spectral measure (recall that  $T_r(t_+, t_-) = U_r^+(t_+)U_r^-(t_-)$ ), hence  $\tilde{T}_r(t_+, t_-)$  sends  $\tilde{\mathcal{M}}_{r,\varphi}$  into itself for  $(t_+, t_-) \in W_R$ . The vector  $\tilde{\Omega}_r$  is cyclic for  $\tilde{\mathcal{M}}_{r,\varphi}$  since  $\tilde{\mathcal{M}}_{r,\varphi}\tilde{\Omega}_r \supset \{(x \otimes \mathbb{1}) \cdot \tilde{R}_\varphi \cdot (\mathbb{1} \otimes y) \cdot \tilde{\Omega}_r\} = \{(x \otimes \mathbb{1}) \cdot (\mathbb{1} \otimes y) \cdot \tilde{\Omega}_r\}$  and the latter is total by the Reeh-Schlieder property of the tensor product net.

We see the separating property of  $\tilde{\Omega}_r$  as follows. Consider a von Neumann algebra

$$\tilde{\mathcal{M}}_{r,\varphi}^1 := \{\text{Ad } \tilde{R}_\varphi(x' \otimes \mathbb{1}), \mathbb{1} \otimes y' : x', y' \in \mathcal{A}_r(W_R)'\}''.$$

One verifies that  $\tilde{\Omega}_r$  is cyclic for  $\tilde{\mathcal{M}}_{r,\varphi}^1$  as above, hence we only have to show that  $\tilde{\mathcal{M}}_{r,\varphi}$  and  $\tilde{\mathcal{M}}_{r,\varphi}^1$  commute. This has been done by Lemma 4.1.  $\square$

One can actually show that  $\tilde{\mathcal{M}}_{r,\varphi}^1 = (\tilde{\mathcal{M}}_{r,\varphi})'$ , so this confusing notation is justified. Indeed, one has only to check the modular group of  $\tilde{\mathcal{M}}_{r,\varphi}$  with respect to  $\tilde{\Omega}_r$  is the same for the tensor product, which follows from the field representation in the next Section and the argument of [23].

In this presentation we took  $\mathcal{A}_r$  as the starting point. It is also possible to take the models in [22] or more general models with more particle spectrum [25]. We will discuss this (slight) generalization in [5]. In this paper we do not consider strict locality of these Borchers triples, although the author expects that a similar proof to [22] should work.

## 5 General arguments for strict locality

We are now concerned with the main problem in the construction of nets through Borchers triples: the strict locality of the models in Section 3. The key argument is the wedge-split property.

In general, if one has an inclusion of von Neumann algebras  $\mathcal{N} \subset \mathcal{M}$ , this is said to be **split** if there is a type I factor  $\mathcal{R}$  such that  $\mathcal{N} \subset \mathcal{R} \subset \mathcal{M}$ . Furthermore, if  $\mathcal{N} \subset \mathcal{M}$  is an inclusion of factors and  $\Omega$  is cyclic and separating for  $\mathcal{M}$ , then the nuclearity of the map

$\mathcal{N} \ni x \mapsto \Delta^{\frac{1}{4}} x \Omega$  implies the split property, where  $\Delta$  is the modular operator for  $\mathcal{M}$  with respect to  $\Omega$  [11, Proposition 2.3].

We say that a Borchers triple  $(\mathcal{M}, T, \Omega)$  is **wedge-split** if  $\text{Ad } T(a)(\mathcal{M}) \subset \mathcal{M}$  is split for any  $a \in W_R$ . Note that  $W_R$  is defined as the *open* wedge and split inclusion for lightlike translation is not required. It is easy to see already from wedge-split property that the intersection  $\mathcal{M} \cap \text{Ad } T(a)(\mathcal{M})$  for  $a \in W_R$  is nontrivial. Let  $\Delta$  be the modular operator for  $\mathcal{M}$  with respect to  $\Omega$ . One says that an inclusion  $\mathcal{N} \subset \mathcal{M}$  satisfies **modular nuclearity** if the map  $\mathcal{N} \ni x \mapsto \Delta^{\frac{1}{4}} x \Omega$  is nuclear. A Borchers triple  $(\mathcal{M}, T, \Omega)$  is said to satisfy modular nuclearity if the inclusion  $\text{Ad } T(\mathcal{M}) \subset \mathcal{M}$  has modular nuclearity for any  $a \in W_R$ . From the above remark, modular nuclearity implies wedge-split property. The strict locality of the models in [22] was proved through modular nuclearity.

Actually, wedge-split inclusion is sufficient for the strict locality [22, Theorem 2.5].

**Theorem 5.1** (Lechner). *If a Borchers triple  $(\mathcal{M}, T, \Omega)$  is wedge-split, then it is strictly local.*

This theorem is stated with the assumption of modular nuclearity, however, the actual proof depends only on wedge-split property.

Let us recall our main construction strategy (Section 3): starting with a given Borchers triple  $(\mathcal{M}_c, T_c, \Omega_c)$  with inner symmetry  $\text{Ad } V_c(\kappa)$  of  $S^1$ , we constructed a new Borchers triple  $(\tilde{\mathcal{M}}_{c,\kappa}, \tilde{T}_c, \tilde{\Omega}_c)$  on the tensor product Hilbert space. As we will see later (Section 6.1), this gives a nontrivial scattering even if the initial triple comes from the free field. This can be generalized to the following program: take a Borchers triple with a good property (either wedge-split property or modular nuclearity) and prove that the construction of Section 3 leads again to Borchers triples with the same property.

We carry out this program in two ways. First we present a proof through wedge-split property and then we use modular nuclearity. Note that strict locality is not “good enough” for this program. We will exhibit examples of strictly local Borchers triples for which the new triples constructed as in Section 3 violate strict locality. Another remark is that our arguments in Section 5.2 through wedge-split property are valid for both  $S^1$ - and  $\mathbb{Z}_N$ -actions, but those in Section 5.3 through modular nuclearity apply so far only to finite cyclic group actions. Already for the simplest compact group  $S^1$  the proofs break down, as we will see.

## 5.1 Common arguments

Let us collect some facts which will be commonly used in the following (see also [33, Section 4.4.1]). In this Subsection, we do not use the properties of Borchers triple.

In the following, we consider only an action of  $S^1$ . The case of  $\mathbb{Z}_N$  is analogous. Let  $\mathcal{M}_c$  be a von Neumann algebra,  $\Omega_c$  a cyclic separating vector for  $\mathcal{M}_c$  and  $V_c(\kappa)$  be a unitary representation of  $S^1$  which preserves  $\Omega_c$  such that  $\alpha_{c,\kappa} := \text{Ad } V_c(\kappa)$  is an automorphism of  $\mathcal{M}_c$  and  $V_c(1) = \mathbb{1}$ . In other words, there is an action  $\alpha_c$  of  $S^1$  on  $\mathcal{M}_c$  which preserves the state  $\langle \Omega_c, \cdot \Omega_c \rangle$ .

## Discrete Fourier expansion on von Neumann algebras

For any element  $x \in \mathcal{M}_c$  and  $l \in \mathbb{Z}$ , we define

$$x_l := \int_0^1 d\kappa \operatorname{Ad} V_c(\kappa)(x) e^{-i2\pi l\kappa},$$

It holds that  $x = \sum_{l \in \mathbb{Z}} x_l$  and  $\operatorname{Ad} V_c(\kappa)(x_l) = \alpha_{c,\kappa}(x_l) = e^{i2\pi l\kappa} x_l$ . The sum is  $*$ -strongly convergent. Let us denote by  $\mathcal{M}_c^{\alpha_c}(l)$  the set of elements  $x$  of  $\mathcal{M}_c$  such that  $\alpha_{c,\kappa}(x) = e^{i2\pi l\kappa} x$ . Then it holds that  $x_l \in \mathcal{M}_c^{\alpha_c}(l)$  and  $\mathcal{M}_c^{\alpha_c}(l_1) \cdot \mathcal{M}_c^{\alpha_c}(l_2) \subset \mathcal{M}_c^{\alpha_c}(l_1 + l_2)$ .

These are well-known facts, however, if necessary the reader is referred to [33, Proposition 4.9] for a proof (for the case of  $\mathbb{Z}_N$ , the proof is easy to adapt).

Next we consider the twisting. As in Section 3, let  $Q_c$  be a self-adjoint operator such that  $V_c(\kappa) = e^{i2\pi\kappa Q_c}$ . Then for  $y \in \mathcal{M}_c^{\alpha_c}(m)$ , it holds that

$$\operatorname{Ad} \tilde{V}_{c,\kappa}(\mathbb{1} \otimes y) = \operatorname{Ad} e^{i2\pi\kappa Q_c \otimes Q_c}(\mathbb{1} \otimes y) = e^{i2\pi m\kappa Q_c} \otimes y = V_c(m\kappa) \otimes y,$$

where  $\tilde{V}_{c,\kappa} = e^{i2\pi\kappa Q_c \otimes Q_c}$  as in Section 3. For the proof, see [33, Lemma 4.10]. The proof is written again for an action of  $S^1$ , however, the adaptation is easy for  $\mathbb{Z}_N$ . Note that one has to consider spectral subspaces of  $V_c$  parametrized by  $\mathbb{Z}_N$ . The identification  $\mathbb{Z}_N = \mathbb{Z}/N\mathbb{Z}$  should be always kept in mind and numbers, e.g.,  $e^{i\frac{2\pi k}{N}}$  are well-defined for  $k \in \mathbb{Z}_N$ .

Finally we remark that  $\tilde{\mathcal{M}}_{c,\kappa}$  is the  $*$ -strong closure of the linear span of the elements of the form  $x_l V_c(m\kappa) \otimes y_m$ , where  $x_l \in \mathcal{M}_c^{\alpha_c}(l)$  and  $y_m \in \mathcal{M}_c^{\alpha_c}(m)$ . Furthermore, any element  $\tilde{x} \in \tilde{\mathcal{M}}_{c,\kappa}$  can be decomposed as follows:

$$\tilde{x} = \sum_{l,m \in \mathbb{Z}} \tilde{x}_{l,m} (V_c(m\kappa) \otimes \mathbb{1}), \quad \tilde{x}_{l,m} \in \mathcal{M}_c \otimes \mathcal{M}_c,$$

and the sum is  $*$ -strongly convergent. This decomposition corresponds to the discrete Fourier expansion with respect to the action  $(\operatorname{Ad} V_c) \otimes (\operatorname{Ad} V_c)$  of the group  $S^1 \times S^1$ . The group  $S^1 \times S^1$  acts also on  $\mathcal{M}_c \otimes \mathcal{M}_c$  and there is a decomposition into components as well. For a corresponding proof, see again [33, Lemma 4.14].

## Modular operator and inner symmetry

Let  $\Delta_c$  and  $J_c$  be the modular operator and the modular conjugation for  $\mathcal{M}_c$  with respect to  $\Omega_c$ . The implementing unitary  $V_c$  of  $\alpha_c$  and  $\Delta_c, J_c$  commute. The modular operator and the modular conjugation for  $\mathcal{M}_c \otimes \mathcal{M}_c$  with respect to  $\tilde{\Omega}_c := \Omega_c \otimes \Omega_c$  are  $\Delta_c \otimes \Delta_c$  and  $J_c \otimes J_c$ , respectively. Now we can determine the modular objects for  $\tilde{\mathcal{M}}_{c,\kappa} = \mathcal{M}_c \otimes \mathbb{1} \vee \operatorname{Ad} \tilde{V}_{c,\kappa}(\mathbb{1} \otimes \mathcal{M}_c)$ .

**Proposition 5.2.** *The operators  $\Delta_c \otimes \Delta_c$  is the modular operator for  $\tilde{\mathcal{M}}_{c,k}$  with respect to  $\tilde{\Omega}_c$ .*

*Proof.* We will give two proofs. The first one is a direct calculation and the second one uses the so-called KMS condition.

A direct calculation goes as follows. We will see that actually the modular conjugation is  $\tilde{V}_{c,\kappa}(J_c \otimes J_c)$ . Since we have already the candidates for the modular objects, we only have to check their actions. First we take  $x_l \in \mathcal{M}_c^{\alpha_c}(l)$  and  $y_m \in \mathcal{M}_c^{\alpha_c}(m)$ . The operator  $x_l V_c(m\kappa) \otimes y_m$  belongs to  $\tilde{\mathcal{M}}_{c,\kappa}$  and it holds that

$$\begin{aligned}
\tilde{V}_{c,\kappa}(J_c \otimes J_c)(\Delta_c \otimes \Delta_c)(x_l V_c(m\kappa) \otimes y_m) \tilde{\Omega}_c &= \tilde{V}_{c,\kappa}(S_c \otimes S_c)(x_l \Omega_c \otimes y_m \Omega_c) \\
&= \tilde{V}_{c,\kappa}(x_l^* \Omega_c \otimes y_m^* \Omega_c) \\
&= e^{i2\pi l m \kappa} (x_l^* \Omega_c \otimes y_m^* \Omega_c) \\
&= (\alpha_{c,-m\kappa}(x_l^*) \Omega_c) \otimes (y_m^* \Omega_c) \\
&= (V_c(-m\kappa) x_l^* \Omega_c) \otimes (y_m^* \Omega_c) \\
&= (x_l V_c(m\kappa) \otimes y_m)^* (\Omega_c \otimes \Omega_c),
\end{aligned}$$

namely, the operator  $\tilde{V}_{c,\kappa}(J_c \otimes J_c)(\Delta_c \otimes \Delta_c)$  acts correctly as the modular involution. Since  $\tilde{\mathcal{M}}_{c,\kappa}$  is the  $*$ -strong closure of the linear span of such elements, the modular involution actually coincides with  $\tilde{V}_{c,\kappa}(J_c \otimes J_c)(\Delta_c \otimes \Delta_c)$ . Then the conclusion follows immediately from the uniqueness of the polar decomposition.

In the second proof, we use the uniqueness of the modular automorphism group with respect to the KMS condition. In general, for a von Neumann algebra  $\mathcal{M}$  and a state  $\psi$ , if there is a one-parameter automorphisms  $\sigma_t$  which satisfies the KMS condition, namely if for any pair  $x, y \in \mathcal{M}$  there is an analytic function  $f(t)$  on the strip  $0 < \Im t < 1$  and continuous on the boundary such that  $f(t) = \psi(\sigma_t(x)y)$  and  $f(t+i) = \psi(y\sigma_t(x))$  for  $t \in \mathbb{R}$ , then  $\sigma_t$  is the modular automorphism [32, Theorem VIII.1.2]. In our case, the state is the vacuum  $\langle \tilde{\Omega}_c, \cdot \tilde{\Omega}_c \rangle$ . This is the KMS state on the von Neumann algebra  $\mathcal{M}_c \otimes \mathcal{M}_c$  with respect to  $\sigma_t^{\tilde{\Omega}_c} = \text{Ad}(\Delta_c^{it} \otimes \Delta_c^{it})$ . Hence for any pair  $\tilde{x}, \tilde{y} \in \mathcal{M}_c \otimes \mathcal{M}_c$  of elements there is an analytic function as above. Actually the decomposition with respect to  $\alpha_c \otimes \alpha_c$  commutes with  $\sigma_t^{\tilde{\Omega}_c}$  (see the remark about discrete Fourier expansion) and it holds that

$$\begin{aligned}
\langle \tilde{\Omega}_c, \sigma_t^{\tilde{\Omega}_c}(\tilde{x}) \tilde{y} \tilde{\Omega}_c \rangle &= \sum_{l,m} \langle \tilde{\Omega}_c, \sigma_t^{\tilde{\Omega}_c}(\tilde{x}_{l,m}) \tilde{y}_{-l,-m} \tilde{\Omega}_c \rangle, \\
\langle \tilde{\Omega}_c, \tilde{y} \sigma_t^{\tilde{\Omega}_c}(\tilde{x}) \tilde{\Omega}_c \rangle &= \sum_{l,m} \langle \tilde{\Omega}_c, \tilde{y}_{-l,-m} \sigma_t^{\tilde{\Omega}_c}(\tilde{x}_{l,m}) \tilde{\Omega}_c \rangle
\end{aligned}$$

by the orthogonality of the vacuum acted on by the component  $\tilde{x}_{l,m}$  etc. Next, by considering the pair  $\tilde{x}_{l,m}, \tilde{y}_{-l,-m}$  and the KMS condition for  $\mathcal{M}_c \otimes \mathcal{M}_c$  there is an analytic function  $\tilde{f}_{l,m}(t) = \langle \tilde{\Omega}_c, \sigma_t^{\tilde{\Omega}_c}(\tilde{x}_{l,m}) \tilde{y}_{-l,-m} \tilde{\Omega}_c \rangle$ ,  $\tilde{f}_{l,m}(t+i) = \langle \tilde{\Omega}_c, \tilde{y}_{-l,-m} \sigma_t^{\tilde{\Omega}_c}(\tilde{x}_{l,m}) \tilde{\Omega}_c \rangle$  (of course the dependence of  $\tilde{f}_{l,m}$  on  $\tilde{x}$  and  $\tilde{y}$  is implicit). Let us turn to  $\tilde{\mathcal{M}}_{c,\kappa}$ . By the same argument as above, an inner product decomposes into suitable combination and the decomposition is compatible with the action of  $\sigma_t^{\tilde{\Omega}_c} = \text{Ad}(\Delta_c^{it} \otimes \Delta_c^{it})$ . For a pair of elements  $\tilde{x}_{l,m}(e^{i2\pi m \kappa} Q_c \otimes \mathbb{1})$

and  $\tilde{y}_{-l,-m}(e^{-i2\pi m\kappa Q_c} \otimes \mathbb{1})$ , where  $\tilde{x}_{l,m}, \tilde{y}_{-l,-m} \in \mathcal{M}_c \otimes \mathcal{M}_c$ , we have

$$\begin{aligned} \langle \tilde{\Omega}_c, \sigma_t^{\tilde{\Omega}_c}(\tilde{x}_{l,m}(e^{i2\pi m\kappa Q_c} \otimes \mathbb{1}))\tilde{y}_{-l,-m}(e^{-i2\pi m\kappa Q_c} \otimes \mathbb{1})\tilde{\Omega}_c \rangle &= e^{-i2\pi lm\kappa} \langle \tilde{\Omega}_c, \sigma_t^{\tilde{\Omega}_c}(\tilde{x}_{l,m})\tilde{y}_{-l,-m}\tilde{\Omega}_c \rangle \\ \langle \tilde{\Omega}_c, \tilde{y}_{-l,-m}(e^{-i2\pi m\kappa Q_c} \otimes \mathbb{1})\sigma_t^{\tilde{\Omega}_c}(\tilde{x}_{l,m}(e^{i2\pi m\kappa Q_c} \otimes \mathbb{1}))\tilde{\Omega}_c \rangle &= e^{-i2\pi lm\kappa} \langle \tilde{\Omega}_c, \tilde{y}_{-l,-m}\sigma_t^{\tilde{\Omega}_c}(\tilde{x}_{l,m})\tilde{\Omega}_c \rangle \end{aligned}$$

One observes that the right hand sides are equal to  $\tilde{f}_{l,m}$  up to the constant  $e^{-i2\pi lm\kappa}$ . In other words, the KMS condition is satisfied for the pair with respect to  $\sigma_t^{\tilde{\Omega}_c}$ . Therefore for an arbitrary linear combination of such components the KMS condition holds as well, thanks to the decomposition of the inner product into  $(l, m)$ -components. Such linear combinations is  $*$ -strongly dense in  $\tilde{\mathcal{M}}_{c,\kappa}$ , hence the above-cited uniqueness theorem applies to see that  $\sigma_t^{\tilde{\Omega}_c} = \text{Ad}(\Delta_c^{it} \otimes \Delta_c^{it})$  is the modular automorphism of  $\tilde{\mathcal{M}}_{c,\kappa}$  with respect to  $\langle \tilde{\Omega}_c, \cdot \tilde{\Omega}_c \rangle$  (indeed, the KMS condition on a  $*$ -strongly dense subalgebra is enough by [9, Proposition 5.7]). One should note that in [9] the KMS condition is defined on a dense set of analytic elements). Since  $\Delta_c \otimes \Delta_c$  preserves the vacuum vector  $\Omega_c \otimes \Omega_c$ , it must coincide with the modular operator.  $\square$

It is interesting to compare our proof with [12, 22] where the modular objects were calculated through unbounded operators affiliated to the von Neumann algebras. One recalls also that in our previous work the modular objects were indirectly determined by scattering theory [33, Section 3].

**Proposition 5.3.** *It holds that  $(\tilde{\mathcal{M}}_{c,\kappa})' = \left( \text{Ad } \tilde{V}_{c,k}(\mathcal{M}'_c \otimes \mathbb{1}) \vee \mathbb{1} \otimes \mathcal{M}'_c \right) =: \tilde{\mathcal{M}}_{c,\kappa}^1$ .*

*Proof.* It can be observed that the right hand side commute with  $\tilde{\mathcal{M}}_{c,\kappa}$  by the same argument as in Theorem 3.3, so we have  $\tilde{\mathcal{M}}_{c,\kappa}^1 \subset (\tilde{\mathcal{M}}_{c,\kappa})'$ . Since we already know that the modular operator for  $(\tilde{\mathcal{M}}_{c,\kappa})'$  with respect to  $\Omega_c \otimes \Omega_c$  is  $\Delta_c^{-1} \otimes \Delta_c^{-1}$  from Proposition 5.2, it is immediate to see that  $\tilde{\mathcal{M}}_{c,\kappa}^1$  is globally invariant under the modular group of  $(\tilde{\mathcal{M}}_{c,\kappa})'$  with respect to  $\Omega_c \otimes \Omega_c$ . Then the two algebras coincide since  $\Omega_c \otimes \Omega_c$  is cyclic for  $\tilde{\mathcal{M}}_{c,\kappa}^1$  by a standard application of Takesaki's theorem [32, Theorem IX.4.2] (see [34, Theorem A.1]).  $\square$

Variations of Propositions 5.2 and 5.3 hold for an action of  $\mathbb{Z}_N$  with trivial changes.

## 5.2 Proof through wedge-split property

Here we take as the starting point a Borchers triple  $(\mathcal{M}_c, T_c, \Omega_c)$  which is wedge-split. As before, we assume that there is an action of  $S^1$  implemented by  $V_c(\kappa)$ . With this action and for  $\kappa \in \mathbb{R}$  we can construct a Borchers triple  $(\tilde{\mathcal{M}}_{c,\kappa}, \tilde{T}_c, \tilde{\Omega}_c)$  as in Section 3. We are going to prove that this is again wedge-split, hence strictly local.

Recall that [15] for a split inclusion  $(\mathcal{N}_c \subset \mathcal{M}_c, \Omega_c)$  equipped with a cyclic separating vector  $\Omega_c$  for  $\mathcal{M}_c, \mathcal{N}_c$  and  $\mathcal{M}_c \cap \mathcal{N}_c$  (where  $\mathcal{N}_c$  will be  $\text{Ad } T_c(a)(\mathcal{M}_c)$  for some  $a \in W_R$ ), there is a *canonical* type I factor  $\mathcal{R}_c$  such that  $\mathcal{N}_c \subset \mathcal{R}_c \subset \mathcal{M}_c$ . Moreover,  $\mathcal{R}_c$  is given by the

formula  $\mathcal{R}_c = \mathcal{N}_c \vee J\mathcal{N}_cJ = \mathcal{M}_c \cap J\mathcal{M}_cJ$ , where  $J$  is the modular conjugation for  $\mathcal{M}_c \cap \mathcal{N}_c$  with respect to  $\Omega_c$ . If  $V_c$  is a unitary operator which preserves  $\Omega_c$  and  $\text{Ad } V_c$  preserves both  $\mathcal{N}_c$  and  $\mathcal{M}_c$ , then  $\mathcal{R}_c$  is preserved under  $\text{Ad } V_c$  as well.

**Lemma 5.4.** *The von Neumann algebra  $\tilde{\mathcal{R}}_{c,\kappa} := \mathcal{R}_c \otimes \mathbb{1} \vee \text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes \mathcal{R}_c)$  is a factor.*

*Proof.* Since we know that  $(\tilde{\mathcal{R}}_{c,\kappa})' = \text{Ad } \tilde{V}_{c,\kappa}(\mathcal{R}'_c \otimes \mathbb{1}) \vee \mathbb{1} \otimes \mathcal{R}'_c$  by Proposition 5.3 (in Section 5.1 we did not assume that the von Neumann algebra comes from a Borchers triple), we only have to show that  $\tilde{\mathcal{R}}_{c,\kappa} \vee (\tilde{\mathcal{R}}_{c,\kappa})' = \mathcal{R}_c \otimes \mathcal{R}'_c \vee \text{Ad } \tilde{V}_{c,\kappa}(\mathcal{R}'_c \otimes \mathcal{R}_c)$  is equal to  $B(\mathcal{H}_c \otimes \mathcal{H}_c)$ .

We show that  $V_c(\kappa) \otimes \mathbb{1}$  and  $\mathbb{1} \otimes V_c(\kappa)$  are contained in  $\tilde{\mathcal{R}}_{c,\kappa} \vee (\tilde{\mathcal{R}}_{c,\kappa})'$  for each  $\kappa \in \mathbb{R}$ . Indeed,  $V_c(\kappa) \otimes \mathbb{1}$  implements an automorphism on  $\mathcal{R}_c \otimes \mathbb{1}$  and since the latter is a type I factor, there is a unitary  $u \in \mathcal{R}_c$  which implements the same automorphism. At the same time  $V_c(\kappa) \otimes \mathbb{1}$  implements an automorphism of  $\mathcal{R}'_c \otimes \mathbb{1}$ , and there is an implementing unitary  $u' \in \mathcal{R}'_c$ . Then  $uu'$  implements  $\text{Ad } V_c(\kappa)$  on  $\mathcal{B}(\mathcal{H}_c) = \mathcal{R}_c \vee \mathcal{R}'_c$ , hence  $uu'V_c(\kappa)^*$  must be a scalar. We may assume  $uu' = V_c(\kappa)$ . Furthermore, obviously  $\text{Ad } V_c(\kappa)(u') = \text{Ad } u'(u') = u'$ , namely  $u'$  is fixed under the automorphism  $\text{Ad } V_c(\kappa)$ , hence  $\text{Ad } \tilde{V}_{c,\kappa}(u' \otimes \mathbb{1}) = u' \otimes \mathbb{1}$ . This implies that  $u' \otimes \mathbb{1} \in (\tilde{\mathcal{R}}_{c,\kappa})'$  hence  $uu' \otimes \mathbb{1} = V_c(\kappa) \otimes \mathbb{1}$  is in  $\tilde{\mathcal{R}}_{c,\kappa} \vee (\tilde{\mathcal{R}}_{c,\kappa})'$ . An analogous proof works for  $\mathbb{1} \otimes V_c(\kappa)$ .

The rest is easy since  $\tilde{V}_{c,\kappa}$  is obtained from the functional calculus of  $V_c(\kappa) \otimes \mathbb{1}$  and  $\mathbb{1} \otimes V_c(\kappa)$ .  $\square$

This Lemma actually works even for an action of  $\mathbb{R}$ . In contrast, we need the periodicity of the action in the following.

**Theorem 5.5.** *The von Neumann algebra  $\tilde{\mathcal{R}}_{c,\kappa} := \mathcal{R}_c \otimes \mathbb{1} \vee \text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes \mathcal{R}_c)$  is a type I factor.*

*Proof.* We have seen that  $\tilde{\mathcal{R}}_{c,\kappa}$  is a factor. What remains is to show that  $\tilde{\mathcal{R}}_{c,\kappa}$  contains a minimal projection. As in Lemma 5.4 we take implementing unitary  $u(\kappa) \in \mathcal{R}_c$  for  $\text{Ad } V_c(\kappa)$ , this time indicating the dependence on  $\kappa$ . By a classical result by Bargmann, we may assume that  $u(\kappa)$  is a one-parameter group of unitaries [2, Section 6a], where our group is  $S^1$  hence there is no nontrivial antisymmetric bilinear form on its Lie algebra  $\mathbb{R}$ .

Thus the spectrum of the representation  $u$  is discrete and contained in  $\mathbb{Z}$ , hence the fixed point  $\mathcal{R}_c^{\alpha_c} = \mathcal{R}_c \cap \{u(\kappa) : \kappa \in \mathbb{R}\}'$  is the relative commutant in  $\mathcal{R}_c$  of the spectral projections of the representation  $u$ . Hence it is the direct sum of type I factors and admits a minimal projection.

Let us take such a minimal projection  $p$  in  $\mathcal{R}_c^{\alpha_c}$ . In general, if  $y \in \mathcal{R}_c^{\alpha_c}$ , then  $\text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes y) = \mathbb{1} \otimes y$  and hence the inclusion  $\mathcal{R}_c \otimes \mathcal{R}_c^{\alpha_c} \subset \tilde{\mathcal{R}}_{c,\kappa}$  holds. Obviously  $p$  must be a subprojection of a discrete point spectrum of  $u$ . Furthermore, it is immediate that  $p \otimes p$  is a minimal projection in  $\mathcal{R}_c \otimes \mathcal{R}_c^{\alpha_c}$ . We claim that it is a minimal projection in  $\tilde{\mathcal{R}}_{c,\kappa}$ . In fact, suppose that  $\tilde{q} \leq p \otimes p$  in  $\tilde{\mathcal{R}}_{c,\kappa}$ . Then  $\tilde{q}$  and  $\mathbb{1} \otimes u(\kappa)$  commute since the spectral projections of the representation  $\mathbb{1} \otimes u$  either contains or disjoint from  $p \otimes p$ , hence so does it hold with  $\tilde{q}$ . This implies that  $\tilde{q}$  belongs to  $\mathcal{R}_c \otimes \mathcal{R}_c^{\alpha_c}$  (consider the discrete Fourier expansion  $\tilde{q} = \sum \tilde{q}_{lm}(V_c(m\kappa) \otimes \mathbb{1})$ ), then it is possible that  $\text{Ad}(1 \otimes u(\kappa))(\tilde{q}) = \tilde{q}$  only if  $\tilde{q}_{lm}$

vanish except  $m = 0$ ), then  $\tilde{q} = p \otimes p$  by the minimality of  $p \otimes p$  in this restricted algebra. This is the minimality of  $p \otimes p$  in  $\tilde{\mathcal{R}}_{c,\kappa}$ .  $\square$

We remark that the intermediate type I factor constructed here is not the canonical one of [15]. An explicit formula for it involves the modular conjugation of the relative commutant, which is only abstractly determined.

The proof of Theorem 5.5 can be easily adapted to the case of an action of  $\mathbb{Z}_N$ .

In terms of wedge-split Borchers triple, we apply Theorem 5.5 with  $\mathcal{N}_c = \text{Ad } T_c(a)(\mathcal{M}_c)$  to see the split inclusion

$$\mathcal{N}_c \otimes \mathbb{1} \vee \text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes \mathcal{N}_c) \subset \mathcal{R}_c \otimes \mathbb{1} \vee \text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes \mathcal{R}_c) \subset \mathcal{M}_c \otimes \mathbb{1} \vee \text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes \mathcal{M}_c),$$

where the middle element is  $\tilde{\mathcal{R}}_{c,\kappa}$  and the last element is  $\tilde{\mathcal{M}}_{c,\kappa}$ . Since  $\tilde{T}_c$  and  $\tilde{V}_{c,\kappa}$  commute,  $\text{Ad } \tilde{T}_c(a)(\tilde{\mathcal{M}}_{c,\kappa}) \subset \tilde{\mathcal{M}}_{c,\kappa}$  is split with an intermediate type I factor  $\tilde{\mathcal{R}}_{c,\kappa}$  (which implicitly depends on  $a$ ). Then we have the following with the help of Theorem 5.1.

**Theorem 5.6.** *If a given Borchers triple  $(\mathcal{M}_c, T_c, \Omega_c)$  with an action of  $S^1$  by inner symmetry is wedge-split, then the Borchers triple  $(\tilde{\mathcal{M}}_{c,\kappa}, \tilde{T}_c, \tilde{\Omega}_c)$  is wedge-split, hence strictly local.*

One can repeat parallel proofs for an action of  $\mathbb{Z}_N$  to obtain the same result.

### 5.3 Proof through modular nuclearity

Let us give another proof of strict locality, based on modular nuclearity. Here we have to restrict ourselves to the case of an action of  $\mathbb{Z}_N$ . Of course one can take an arbitrary finite subgroup  $\mathbb{Z}_N$  in  $S^1$ , hence physically this should not be considered as an essential restriction. Let  $(\mathcal{M}_{c,k}, \tilde{T}_c, \tilde{\Omega}_c)$  again be constructed as in Section 3 for a fixed  $k \in \mathbb{Z}_N$ . Again, thanks to Theorem 5.1, it is enough to prove the modular nuclearity of the new triple.

Let us start with a trivial observation.

**Lemma 5.7.** *Let  $(\mathcal{M}_c, T_c, \Omega_c)$  be a Borchers triple with modular nuclearity. Then the triple  $(\mathcal{M}_c \otimes \mathcal{M}_c, \tilde{T}_c, \tilde{\Omega}_c)$  has modular nuclearity.*

*Proof.* The modular objects are the tensor products as well, hence the nuclearity norm simply gets squared.  $\square$

**Theorem 5.8.** *Let  $(\mathcal{M}_c, T_c, \Omega_c)$  be a Borchers triple with modular nuclearity. Then the triple  $(\tilde{\mathcal{M}}_{c,k}, \tilde{T}_c, \tilde{\Omega}_c)$  has modular nuclearity.*

*Proof.* For a fixed  $a \in W_R$ , we have to show that the inclusion  $\text{Ad } \tilde{T}_c(a)(\tilde{\mathcal{M}}_{c,k}) \subset \tilde{\mathcal{M}}_{c,k}$  has modular nuclearity with respect to  $\tilde{\Omega}_c$ . By assumption and Lemma 5.7 we know that  $(\text{Ad } \tilde{T}_c(a)(\mathcal{M}_c \otimes \mathcal{M}_c) \subset \mathcal{M}_c \otimes \mathcal{M}_c)$  has modular nuclearity. This means that the map

$$\mathcal{M}_c \otimes \mathcal{M}_c \ni \tilde{x} \longmapsto \left( \Delta_c^{\frac{1}{4}} \otimes \Delta_c^{\frac{1}{4}} \right) \tilde{x}(\Omega_c \otimes \Omega_c)$$

is a nuclear map.

Let us consider an element  $\tilde{x}$  of  $\mathcal{M}_c \otimes \mathcal{M}_c$ . As we have seen in the end of Section 5.1, it can be decomposed as  $\tilde{x} = \sum_{l,m} \tilde{x}_{l,m}$ , and  $\tilde{x}_{l,m}$  are the matrix components in  $\mathcal{M}_c \otimes \mathcal{M}_c$ . For such  $\tilde{x}$ , we define an element  $\tau_k(\tilde{x}) \in \tilde{\mathcal{M}}_{c,k}$  as follows:

$$\tau_k(\tilde{x}) = \sum_{l,m} \tilde{x}_{l,m} (V_c^{km} \otimes \mathbb{1}).$$

This is a finite sum, thus there is no problem of convergence. This map  $\tau_k$  is onto, as any element in  $\tilde{\mathcal{M}}_{c,k}$  can be decomposed in the form above. It is important to observe that the action on the vector  $\tilde{\Omega}_c$  is unchanged. In other words, it holds that  $\tilde{x}\tilde{\Omega}_c = \tau_k(\tilde{x})\tilde{\Omega}_c$ .

Now, we know that the modular operator of  $\tilde{\mathcal{M}}_{c,k}$  with respect to  $\tilde{\Omega}_c$  is  $\Delta_c \otimes \Delta_c$ , hence the map in question is  $\tau_k(\tilde{x}) \mapsto (\Delta_c^{\frac{1}{4}} \otimes \Delta_c^{\frac{1}{4}}) \tau_k(\tilde{x}) \tilde{\Omega}_c = (\Delta_c^{\frac{1}{4}} \otimes \Delta_c^{\frac{1}{4}}) \tilde{x} \tilde{\Omega}_c$ . By assumption and Lemma 5.7 we know that the map  $\tilde{x} \mapsto (\Delta_c^{\frac{1}{4}} \otimes \Delta_c^{\frac{1}{4}}) \tau_k(\tilde{x}) \tilde{\Omega}_c$  is nuclear.

$$\begin{array}{ccccc} \tilde{x} & \xrightarrow{\quad} & \tilde{x}\tilde{\Omega}_c = \tau_k(\tilde{x})\tilde{\Omega}_c & \xrightarrow{\quad} & (\Delta_c^{\frac{1}{4}} \otimes \Delta_c^{\frac{1}{4}}) \tilde{x}\tilde{\Omega}_c \\ \downarrow & & \nearrow & & \\ \tau_k(\tilde{x}) & & & & \end{array}$$

In the diagram above, the left triangle commutes and the straight line above is nuclear. Since a composition of a bounded linear map and a nuclear map is again nuclear, we only have to show that  $\tau_k^{-1}$  is bounded. For this purpose, let us recall how explicitly  $\{\tilde{x}_{l,m}\}$  are obtained (see [33, Proposition 4.9] for a corresponding formula for  $S^1$ -action):

$$\tilde{x}_{l,m} = \frac{1}{N^2} \sum_{j_1, j_2} e^{-i \frac{2\pi(j_1 l + j_2 m)}{N}} \text{Ad}(V_c^{j_1} \otimes V_c^{j_2})(\tilde{x}).$$

Correspondingly for the decomposition of  $\tau_k(\tilde{x})$ , we have

$$\tilde{x}_{l,m} (V_c^{km} \otimes \mathbb{1}) = \frac{1}{N^2} \sum_{j_1, j_2} e^{-i \frac{2\pi(j_1 l + j_2 m)}{N}} \text{Ad}(V_c^{j_1} \otimes V_c^{j_2})(\tau_k(\tilde{x})).$$

In particular, we see  $\|\tilde{x}_{l,m}\| \leq \|\tau_k(\tilde{x})\|$  and hence  $\|\tilde{x}\| = \|\sum_{l,m} \tilde{x}_{l,m}\| \leq N^2 \|\tau_k(\tilde{x})\|$ . In other words,  $\tau_k^{-1}$  is bounded by  $N^2$ .  $\square$

Note that here the finiteness of  $N$  is crucial. The author does not know if the same holds for the action of e.g.  $S^1$ , although one can always take an arbitrary finite subgroup  $\mathbb{Z}_N$  of  $S^1$ .

## 5.4 Counterexamples from massless case

In previous Sections we started with a strictly local Borchers triple, constructed new triples and proved strict locality. Although our method may seem quite simple, it is neither trivial nor purely group-theoretic. In order to see this, we take the massless models, for which one can construct Borchers triples but strict locality fails. See also [33, Section 4.4].

Let  $(\mathcal{A}_0, T_0, \Omega_0)$  be any strongly additive conformal (diffeomorphism covariant) net with an action of  $\mathbb{Z}_N$  by inner symmetry (there are many such nets, e.g. the tensor product of the  $U(1)$ -current or the loop group nets). A two-dimensional massless net can be constructed by tensor product:  $\mathcal{A}(I_+ \times I_-) := \mathcal{A}_0(I_+) \otimes \mathcal{A}_0(I_-)$ ,  $T(t_+, t_-) := T_0(t_+) \otimes T_0(t_-)$ ,  $\Omega := \Omega_0 \otimes \Omega_0$ . On the net  $\mathcal{A}$  there is an action of  $\mathbb{Z}_N$  by inner symmetry which acts only on the left component of  $\mathcal{A}_0(I_+) \otimes \mathcal{A}_0(I_-)$ . Let  $Q$  be the generator with  $\text{sp } Q \subset \mathbb{Z}$ , which is of the form  $Q_0 \otimes \mathbb{1}$ . With the wedge algebra  $\mathcal{M} := \mathcal{A}_0(\mathbb{R}_-) \otimes \mathcal{A}_0(\mathbb{R}_+)$ , the twisted Borchers triple is given by

$$\begin{aligned} \tilde{\mathcal{M}}_{c,k} &:= \mathcal{M} \otimes \mathbb{1} \vee \text{Ad } e^{i\frac{2\pi k}{N}Q \otimes Q}(\mathbb{1} \otimes \mathcal{M}) \\ &= \mathcal{A}_0(\mathbb{R}_-) \otimes \mathcal{A}_0(\mathbb{R}_+) \otimes \mathbb{1} \vee \text{Ad } e^{i\frac{2\pi k}{N}Q \otimes Q}(\mathbb{1} \otimes \mathcal{M}) \\ &\cong \left( \mathcal{A}_0(\mathbb{R}_-) \otimes \mathbb{1} \vee \text{Ad } e^{i\frac{2\pi k}{N}Q_0 \otimes Q_0}(\mathbb{1} \otimes \mathcal{A}_0(\mathbb{R}_-)) \right) \otimes (\mathcal{A}_0(\mathbb{R}_+) \otimes \mathcal{A}_0(\mathbb{R}_+)), \end{aligned}$$

where in the last line we interchanged the second and the third components for brevity. We consider the relative commutant of the wedge shifted by  $a = (a_+, a_-)$ ,  $a_+ < 0$ ,  $a_- > 0$ . With the above interchanged notation, it is clear that the  $--$ -component (the 3rd and 4th tensor components) of the intersection is simply  $\mathcal{A}_0((0, a_-)) \otimes \mathcal{A}_0((0, a_-))$ . As for the  $+-$  component (the 1st and 2nd tensor components), one observes that it is almost same as the intersection calculated in [33, Theorem 4.16]. The only change is that the direction of the 2nd component is reversed. This does not affect the proof, indeed, the inner symmetry commutes with translation and the positivity of energy is used only through Reeh-Schlieder property hence is not essential. Thus we have

$$\tilde{\mathcal{M}}_{c,k} \cap \text{Ad } \tilde{T}_c(a)(\tilde{\mathcal{M}}_{c,k}) = (\mathcal{A}_0^{\alpha_0}((a_+, 0)) \otimes \mathcal{A}_0^{\alpha_0}((a_+, 0))) \otimes (\mathcal{A}_0((0, a_-)) \otimes \mathcal{A}_0((0, a_-))),$$

where  $\mathcal{A}_0^{\alpha_0}$  denotes the fixed point with respect to  $\text{Ad } e^{i\frac{2\pi k}{N}Q_0}$ . In particular with  $k = 1$ , this does not satisfy the Reeh-Schlieder property and strict locality fails if  $\mathbb{Z}_N$  acts nontrivially.

This counterexample shows that our proof of locality is by no means purely group-theoretic. Namely, in order to obtain strict locality, it is necessary to assume stronger property than strict locality itself of the original net (wedge-split property or modular nuclearity as above). The above massless counterexample appears to be related to the subtlety in massless bootstrap program: the convergence of form factors is typically worse in massless models and even the fundamental commutativity theorem relies on the behavior of form factors, which is also worse (c.f. [31]). Hence if one aims at constructing Wightman fields or Haag-Kastler net out of form factors, the problem of convergence is inevitable.

## 6 Realization as deformed fields and scattering theory

In Sections 3 and 4 we constructed families of Borchers triples more or less operator-algebraically. Such a construction was also useful for the argument of strict locality for one case (Section 5). However, we still have to show that the resulting nets of von Neumann algebras are really new, or more desirably have nontrivial interaction and for this purpose the previous presentation is not very convenient. Fortunately, it turns out that the nets are accompanied by (wedge-local) quantum fields which create one-particle states and the scattering process can be calculated [8]. We follow the notations of [25]. The author expects that this field presentation would help to prove strict locality for the construction in Section 4.

### 6.1 The complex free field and inner symmetry

#### The Zamolodchikov-Fadeev algebra

Here we consider the construction in Section 3 applied to the complex free massive field net. The complex massive free net is given simply by the tensor product  $\mathcal{M}_c := \mathcal{M}_r \otimes \mathcal{M}_r$ ,  $T_c := T_r \otimes T_r$ ,  $\Omega_c := \Omega_r \otimes \Omega_r$ . The Hilbert space  $\mathcal{H}_c := \mathcal{H}_r \otimes \mathcal{H}_r$  is canonically isomorphic to the Fock space  $\mathcal{F}(\mathcal{H}_1 \oplus \mathcal{H}_1)$ . The  $U(1)$  symmetry transformation of the complex field is constructed as follows: On the “one-particle space”  $\mathcal{H}_1 \oplus \mathcal{H}_1$  we consider the following operator

$$V_1(\kappa) \begin{pmatrix} \xi \\ \eta \end{pmatrix} = \begin{pmatrix} \cos 2\pi\kappa & -\sin 2\pi\kappa \\ \sin 2\pi\kappa & \cos 2\pi\kappa \end{pmatrix} \begin{pmatrix} \xi \\ \eta \end{pmatrix}, \quad \kappa \in \mathbb{R}$$

The second quantized promotion to  $\mathcal{H}_c$  is denoted by  $V_c(\kappa) := \Gamma(V_1(\kappa))$ . The operator  $V_c(\kappa)$  obviously commutes with  $T_c$  and preserves  $\Omega_c$ . Moreover, for field operators one has

$$\begin{aligned} \text{Ad } V_c(\kappa)(\phi(f) \otimes \mathbb{1}) &= \cos 2\pi\kappa(\phi(f) \otimes \mathbb{1}) + \sin 2\pi\kappa(\mathbb{1} \otimes \phi(f)), \\ \text{Ad } V_c(\kappa)(\mathbb{1} \otimes \phi(g)) &= -\sin 2\pi\kappa(\phi(g) \otimes \mathbb{1}) + \cos 2\pi\kappa(\mathbb{1} \otimes \phi(g)), \end{aligned}$$

and it holds that  $\text{Ad } V_c(\kappa)(e^{i(\phi(f) \otimes \mathbb{1})}) = e^{i \cos 2\pi\kappa \phi(f)} \otimes e^{i \sin 2\pi\kappa \phi(f)}$  and  $\text{Ad } V_c(\kappa)(e^{i(\mathbb{1} \otimes \phi(g))}) = e^{-i \sin 2\pi\kappa \phi(g)} \otimes e^{i \cos 2\pi\kappa \phi(g)}$ . By considering  $f$  and  $g$  supported in  $W_R$ , we conclude that  $\text{Ad } V_c(\kappa)(\mathcal{M}_c) = \mathcal{M}_c$ . In other words,  $V_c(\kappa)$  implements an inner symmetry of the group  $\mathbb{R}/\mathbb{Z} \cong S^1$ .

As we have seen in Section 5, our proof of strict locality works for an action of  $S^1$ . Hence in the following we consider only that case.

Since  $(\mathcal{M}_r, T_r, \Omega_r)$  is wedge-split, so is the tensor product  $(\mathcal{M}_c, T_c, \Omega_c)$ . By considering the above action of  $S^1$  by inner symmetry, one can construct Borchers triples  $(\tilde{\mathcal{M}}_{c,\kappa}, \tilde{T}_c, \tilde{\Omega}_c)$  as in Section 3, which we know to be strictly local by Section 5.

We first take a closer look at the action of  $V_c(\kappa)$ . The matrix  $V_1(\kappa)$  expressed above can be diagonalized by  $\frac{1}{\sqrt{2}} \begin{pmatrix} \mathbb{1} & i\mathbb{1} \\ \mathbb{1} & -i\mathbb{1} \end{pmatrix}$  into  $\begin{pmatrix} e^{i2\pi\kappa} & 0 \\ 0 & e^{-i2\pi\kappa} \end{pmatrix}$ . Correspondingly we define  $\mathcal{H}_{1,\pm} := \{\psi \oplus \pm i\psi : \psi \in \mathcal{H}_1\}$ . Then the full Fock space  $\mathcal{H}_c$  can be decomposed into

$\mathbb{Z}$ -graded subspaces  $\mathcal{H}_c = \bigoplus_{l \in \mathbb{Z}} \mathcal{H}_c^l$  and we may assume that the generator  $Q_c$  of  $V_c(\kappa)$  acts by  $l\mathbb{1}$  on  $\mathcal{H}_c^l$  (by definition of the grading), and  $V_c(\kappa) = e^{i2\pi\kappa Q_c}$ . Hence on the Hilbert space of our interest  $\tilde{\mathcal{H}}_c = \mathcal{H}_c \otimes \mathcal{H}_c = \bigoplus \mathcal{H}_c^l \otimes \mathcal{H}_c^m$ , it is clear that  $Q_c \otimes Q_c$  acts by  $lm\mathbb{1}$  on  $\mathcal{H}_c^l \otimes \mathcal{H}_c^m$ .

Now the operator  $a_+^\dagger(\psi) := a^\dagger(\psi \oplus (i\psi))$  on  $\mathcal{H}_c$  increments the grading and so does  $a_-(\psi) := a(\psi \oplus (-i\psi))$ . On the other hand,  $a_-^\dagger(\psi) := a^\dagger(\psi \oplus (-i\psi))$  and  $a_+(\psi) := a(\psi \oplus (i\psi))$  decrement the grading.

Now it is easy to see the following twisted commutation relations:

$$\begin{aligned} a_\pm^\dagger(\psi_1) \otimes \mathbb{1} \cdot \text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes a_\pm^\dagger(\psi_2)) - e^{\pm(\mp i2\pi\kappa)} \cdot \text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes a_\pm^\dagger(\psi_2)) \cdot a_\pm^\dagger(\psi_1) \otimes \mathbb{1} &= 0, \\ a_\pm(\psi_1) \otimes \mathbb{1} \cdot \text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes a_\pm^\dagger(\psi_2)) - e^{\mp(\mp i2\pi\kappa)} \cdot \text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes a_\pm^\dagger(\psi_2)) \cdot a_\pm(\psi_1) \otimes \mathbb{1} &= 0, \\ a_\pm^\dagger(\psi_1) \otimes \mathbb{1} \cdot \text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes a_\pm(\psi_2)) - e^{\pm(\pm i2\pi\kappa)} \cdot \text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes a_\pm(\psi_2)) \cdot a_\pm^\dagger(\psi_1) \otimes \mathbb{1} &= 0, \\ a_\pm(\psi_1) \otimes \mathbb{1} \cdot \text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes a_\pm(\psi_2)) - e^{\mp(\pm i2\pi\kappa)} \cdot \text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes a_\pm(\psi_2)) \cdot a_\pm(\psi_1) \otimes \mathbb{1} &= 0, \end{aligned}$$

where the signs  $\pm$  etc. in the first term correspond to respectively to  $\pm$  etc. in the constant factor in the second term. The commutation relation between objects with or without  $\text{Ad } \tilde{V}_{c,\kappa}$  follows trivially from the usual ones. Namely, we have

$$[a_\pm^\dagger(\psi_1), a_\pm^\dagger(\psi_2)] = 0, \quad [a_\pm^\dagger(\psi_1), a_\pm(\psi_2)] = \langle \psi_2, \psi_1 \rangle \mathbb{1},$$

and all other combinations commute (note that  $\langle \cdot, \cdot \rangle$  is linear in the second argument and  $a^\dagger(\cdot)$  is linear and  $a(\cdot)$  is antilinear). In other words, these operator-valued distributions satisfy the Zamolodchikov-Fadeev algebra, with the S-matrix given by the phase factors. This two-particle scattering matrix (see below) does not depend on the rapidity  $\theta = \log p$ .

Note that this set of commutation relations can be summarized in the form of matrix. We take a basis  $\{e_{1,+}, e_{1,-}, e_{2,+}, e_{2,-}\}$  on  $\mathbb{C}^2 \otimes \mathbb{C}^2$  and accordingly  $\{e_{1,+} \otimes e_{1,+}, e_{1,+} \otimes e_{1,-}, e_{1,-} \otimes e_{1,+}, e_{1,-} \otimes e_{1,-}, e_{2,+} \otimes e_{2,+}, e_{2,+} \otimes e_{2,-}, \dots\}$  on  $(\mathbb{C}^2 \otimes \mathbb{C}^2) \otimes (\mathbb{C}^2 \otimes \mathbb{C}^2)$ , where

$$(\mathcal{H}_r \oplus \mathcal{H}_r) \oplus (\mathcal{H}_r \oplus \mathcal{H}_r) = (\mathcal{H}_r \otimes \mathbb{C}^2) \otimes \mathbb{C}^2$$

is understood. The signs  $\pm$  refer to the structure of the complex free field, while indices 1, 2 are the first and the second copies of the field. The two-particle S-matrix  $\tilde{S}_{c,\kappa}(\theta)$  is given on this basis by (note that this is constant with respect to  $\theta$  and the dependence on

the choice of  $N$  is always implicitly understood):

$$\begin{pmatrix} 1 & & & \\ & 1 & & \\ & & e^{i2\pi\kappa} & \\ & & & e^{-i2\pi\kappa} \\ \hline & 1 & & \\ & & 1 & \\ & & & e^{-i2\pi\kappa} \\ & & & e^{i2\pi\kappa} \\ \hline e^{-i2\pi\kappa} & & & \\ & e^{i2\pi\kappa} & & \\ & & 1 & \\ & & & 1 \\ \hline e^{i2\pi\kappa} & & & \\ & & e^{-i2\pi\kappa} & \\ & & & 1 \\ & & & & 1 \end{pmatrix}$$

And it is straightforward that this complies the conditions of [25, Definition 2.1] if the charge conjugation  $J_c$  is introduced which exchanges  $+$  and  $-$  fields in each component. This is said to be diagonal in the sense of [25, Section 6]. The advantage of our methods is that the strict locality can be seen as an immediate consequence of Section 5.

This two-particle S-matrix is nontrivial only between different components. One could say that the interaction occurs only between particles of different species but there is no self-interaction. This is clear also from the construction: one component remains unchanged and the other component is just shifted by a unitary equivalence, thus the twisting exists only between different components.

With the help of the analysis [25], we have the following.

**Theorem 6.1.** *The triple  $(\tilde{\mathcal{M}}_{c,\kappa}, \tilde{T}_c, \tilde{\Omega}_c)$  is strictly local and the corresponding Haag-Kastler net is asymptotically complete and interacting and the S-matrix is factorizing and its two-particle S-matrix is given as above.*

### Comparison of von Neumann algebras

By definition we have  $\tilde{\mathcal{M}}_{c,\kappa} = \mathcal{M}_c \otimes \mathbb{1} \vee \text{Ad } \tilde{V}_{c,\kappa}(\mathbb{1} \otimes \mathcal{M}_c)$  and  $\mathcal{M}_c$  is generated by the exponential of fields  $\phi(f) \otimes \mathbb{1}, \mathbb{1} \otimes \phi(g)$ , where  $f, g$  are real test functions with  $\text{supp } f, \text{supp } g \subset W_R$  and  $\mathcal{H}_c = \mathcal{H}_r \otimes \mathcal{H}_r$  is understood. First let us consider the  $\mathcal{M}_c \otimes \mathbb{1}$  component. Following [25], we consider pairs of complex valued test functions  $f, g$  such that  $\bar{f} = g$ . The complex

field which generates the wedge algebra in [25] is given in our notation by

$$\begin{aligned}
\phi_c(f \oplus g) &= a_c^\dagger(f^+ \oplus g^+) + a_c(J_1(f^- \oplus g^-)) \\
&:= a_+^\dagger(f^+) + a_-^\dagger(g^+) + a_+(\overline{g^-}) + a_-(\overline{f^-}) \\
&= \phi(f) \otimes \mathbb{1} + \mathbb{1} \otimes i\phi(f) + \phi(g) \otimes \mathbb{1} + \mathbb{1} \otimes i\phi(g) \\
&= \phi(f + \overline{f}) \otimes \mathbb{1} + \mathbb{1} \otimes i\phi(f - \overline{f}) \\
&= \phi(2\Re f) \otimes \mathbb{1} - \mathbb{1} \otimes \phi(2\Im f),
\end{aligned}$$

where  $J_1(\xi \oplus \eta) = \overline{\eta} \oplus \overline{\xi}$  and  $\xi, \eta \in L^2(\mathbb{R}, d\theta)$  and  $f^\pm(\theta), g^\pm(\theta)$  are defined as before. Namely, the fields in complex and real basis are just the linear combination of each other. Note that  $\phi$  and  $a^\dagger$  are linear but  $a$  is antilinear.

Similarly, one has  $\text{Ad } \widetilde{V}_{c,\kappa}(\phi_c(f \oplus g)) = \text{Ad } \widetilde{V}_{c,\kappa}(\phi_c(2\Re f) \otimes \mathbb{1} - \mathbb{1} \otimes \phi_c(2\Im f))$ . From this one easily shows that the wedge algebra generated by the Zamolodchikov-Fadeev fields in the sense of [25] is equal to  $\widetilde{\mathcal{M}}_{c,\kappa} = \mathcal{M}_c \otimes \mathbb{1} \vee \text{Ad } \widetilde{V}_{c,\kappa}(\mathbb{1} \otimes \mathcal{M}_c)$ .

## Relation to the Federbush model

One notices that this S-matrix is very similar to the one of the Federbush model [28, 14, 30], although here the fields are bosonic. However, our procedure can be easily adapted to fermionic nets. Moreover, in the traditional approach there were technical problems: One can construct local fields only for small coupling constant [29], or if one takes the bootstrap approach, the convergence of form factors are not clear [1]. Here this problem is completely solved. We can prove the existence of local operators for any value of  $\kappa$  if we consider the action of  $S^1$ . By comparing the S-matrix, this corresponds to an arbitrary value of the coupling constant.

More importantly, our construction is not restricted to the Federbush models. One can take any wedge-split net with inner symmetry. This contains, for example, the tensor product of one of Lechner's models [22], instead of the real free field. One can consider  $n$  copies of the real free field, which have  $O(n)$  symmetry, then take any subgroup of  $O(n)$  isomorphic to  $S^1$  or  $\mathbb{Z}_N$ . This should correspond to the Lie-algebraic generalization of the Federbush models, whose form factors were proposed in [14]. It works also with  $n$  copies of one of Lechner's models. Furthermore, the constructed net admits again inner symmetry and is wedge-split, hence one can repeat the construction to obtain further new models (on a bigger Hilbert space).

## Some technical remarks on inner symmetry

The wedge-algebra of our two-dimensional nets is given by the tensor product twisted by the inner symmetry. One realizes that the whole net still admits an action of  $S^1 \times S^1$  by inner symmetry and the fixed point is the same for the untwisted tensor product. In other words, the fixed point net has different (free and interacting) extension. This does not contradict the DHR reconstruction theorem [16, 17], which does not work with a compact group in two dimensions.

Our models violate the parity symmetry, which is clear from the S-matrix. However, the extended parity which interchanges the two components is preserved (see [30, Section 6.3.3]). As noted in [10, Theorem 3.3], the parity symmetry is essential for the Bisognano-Wichmann property in two dimensions. Although this is not necessarily related to our models, we present a simple counterexample. One takes the complex free field, which admits  $S^1$ -inner symmetry with Bisognano-Wichmann property. Then one can simply replace the Lorentz boosts by the composition of Lorentz boosts and the inner symmetry. This still satisfies all the axioms of net but violates the parity symmetry which must have the appropriate commutation relation with the boosts. Accordingly, the inner symmetry and Poincaré symmetry do not necessarily commute. For example, we can take a net with a noncommutative Lie group symmetry and replace the boosts as above. The proof in four dimensions [15, Theorem 10.4] does not work in two dimensions since the Lorentz group is abelian, and hence the Poincaré group has finite dimensional unitary representations.

## 6.2 Real free field and Longo-Witten endomorphisms

Let us see that the Borchers triples constructed in Section 4 admit a wedge-local field interpretation. Since the twisting operator  $\tilde{R}_\varphi$  is given on each particle number space, the calculation is straightforward. We only need the following commutation relations:

$$a^\dagger(\psi_1) \otimes \mathbb{1} \cdot \text{Ad } \tilde{R}_\varphi(\mathbb{1} \otimes a^\dagger(\psi_2)) - \int dp dp' \psi_1(p) \psi_2(p') \varphi\left(\frac{p}{p'}\right) \text{Ad } \tilde{R}_\varphi(\mathbb{1} \otimes a^\dagger(p')) \cdot a^\dagger(p) \otimes \mathbb{1} = 0,$$

or with rapidity  $\theta = \log p$  and in terms of operator-valued distributions one has

$$a^\dagger(\theta) \otimes \mathbb{1} \cdot \text{Ad } \tilde{R}_\varphi(\mathbb{1} \otimes a^\dagger(\theta')) - \varphi(e^{\theta-\theta'}) \text{Ad } \tilde{R}_\varphi(\mathbb{1} \otimes a^\dagger(\theta')) \cdot a^\dagger(\theta) \otimes \mathbb{1} = 0.$$

Note that if  $\varphi$  is an inner symmetric function, then  $\varphi(e^\theta)$  is a bounded analytic function in the strip  $0 < \Im \theta < \pi$  and  $\varphi(e^{i\pi-\theta}) = \varphi(-e^{-\theta})$  has the same property. In the matrix form, it can be written as

$$\tilde{S}_{r,\varphi}(\theta) = \begin{pmatrix} 1 & & \\ & \varphi(e^\theta) & \\ & \varphi(-e^{-\theta}) & \\ & & 1 \end{pmatrix}$$

on the basis  $\{e_1 \otimes e_1, e_1 \otimes e_2, e_2 \otimes e_1, e_2 \otimes e_2\}$ . This is again a diagonal S-matrix. From its simple form, it is expected that the proof of modular nuclearity is similar to the one in [22].

It is clear that also in these models the interaction occurs only between different components.

## 7 Summary and outlook

In this paper, we presented a novel procedure to obtain interacting quantum field models realized as nets of observables. One can view this procedure as first preparing a pair of models then making them couple.

The method is thoroughly operator-algebraic and in the most abstract setting of Section 5 no field picture (wedge-local or not) is required. On the other hand, the interaction is fairly weak. When the field description is available, a pair of particles of different species obtains a phase during the interaction and no momentum transfer occurs. A connection with wedge-local field approach and our previously constructed Borchers triples has been found in [24]. It is worth investigating how to obtain more general integrable models (e.g. [22, 25]) purely operator-algebraically, where a “self-interaction” occurs.

Furthermore, an interesting variant has been obtained in [6]. Strict locality of the models therein is not known, but S-matrix shows a phenomenon which resembles particle production. It is desired to establish strict locality of these models or to find massive correspondents.

More ambitiously, certain relations are claimed between integrable models and higher-dimensional gauge theories (e.g. [4]). The author wishes to study such connections with analytical approach.

### Acknowledgment.

I would like to thank Marcel Bischoff and Karl-Henning Rehren for helpful discussions.

## References

- [1] H. Babujian and M. Karowski. Towards the construction of Wightman functions of integrable quantum field theories. In *Proceedings of 6th International Workshop on Conformal Field Theory and Integrable Models*, volume 19, pages 34–49, 2004.
- [2] V. Bargmann. On unitary ray representations of continuous groups. *Ann. of Math. (2)*, 59:1–46, 1954.
- [3] Hellmut Baumgärtel. *Operator algebraic methods in quantum field theory*. Akademie Verlag, Berlin, 1995.
- [4] Niklas Beisert, Changrim Ahn, Luis F. Alday, and et al. Review of AdS/CFT integrability: An overview. *Lett. Math. Phys.*, 99(1-3):3–32, 2012.
- [5] Marcel Bischoff and Yoh Tanimoto. Integrable QFT and Longo-Witten endomorphisms. *in preparation.*, 2012.
- [6] Marcel Bischoff and Yoh Tanimoto. Construction of Wedge-Local Nets of Observables through Longo-Witten Endomorphisms. II. *Comm. Math. Phys.*, 317(3):667–695, 2013.

- [7] H.-J. Borchers. The CPT-theorem in two-dimensional theories of local observables. *Comm. Math. Phys.*, 143(2):315–332, 1992.
- [8] Hans-Jürgen Borchers, Detlev Buchholz, and Bert Schroer. Polarization-free generators and the  $S$ -matrix. *Comm. Math. Phys.*, 219(1):125–140, 2001.
- [9] Ola Bratteli and Derek W. Robinson. *Operator algebras and quantum statistical mechanics. 2. Equilibrium states. Models in quantum statistical mechanics.* Texts and Monographs in Physics. Springer-Verlag, Berlin, second edition, 1997.
- [10] R. Brunetti, D. Guido, and R. Longo. Modular structure and duality in conformal quantum field theory. *Comm. Math. Phys.*, 156(1):201–219, 1993.
- [11] Detlev Buchholz, Claudio D’Antoni, and Roberto Longo. Nuclear maps and modular structures. I. General properties. *J. Funct. Anal.*, 88(2):233–250, 1990.
- [12] Detlev Buchholz and Gandalf Lechner. Modular nuclearity and localization. *Ann. Henri Poincaré*, 5(6):1065–1080, 2004.
- [13] Detlev Buchholz, Gandalf Lechner, and Stephen J. Summers. Warped convolutions, Rieffel deformations and the construction of quantum field theories. *Comm. Math. Phys.*, 304(1):95–123, 2011.
- [14] O. A. Castro-Alvaredo and A. Fring. Form factors from free fermionic Fock fields, the Federbush model. *Nuclear Phys. B*, 618(3):437–464, 2001.
- [15] S. Doplicher and R. Longo. Standard and split inclusions of von Neumann algebras. *Invent. Math.*, 75(3):493–536, 1984.
- [16] Sergio Doplicher and John E. Roberts. A new duality theory for compact groups. *Invent. Math.*, 98(1):157–218, 1989.
- [17] Sergio Doplicher and John E. Roberts. Why there is a field algebra with a compact gauge group describing the superselection structure in particle physics. *Comm. Math. Phys.*, 131(1):51–107, 1990.
- [18] Wojciech Dybalski and Yoh Tanimoto. Asymptotic completeness in a class of massless relativistic quantum field theories. *Comm. Math. Phys.*, 305(2):427–440, 2011.
- [19] Fabrizio Gabbiani and Jürg Fröhlich. Operator algebras and conformal field theory. *Comm. Math. Phys.*, 155(3):569–640, 1993.
- [20] Rudolf Haag. *Local quantum physics.* Texts and Monographs in Physics. Springer-Verlag, Berlin, second edition, 1996. Fields, particles, algebras.
- [21] Gandalf Lechner. Polarization-free quantum fields and interaction. *Lett. Math. Phys.*, 64(2):137–154, 2003.

- [22] Gandalf Lechner. Construction of quantum field theories with factorizing  $S$ -matrices. *Comm. Math. Phys.*, 277(3):821–860, 2008.
- [23] Gandalf Lechner. Deformations of quantum field theories and integrable models. *Comm. Math. Phys.*, 312(1):265–302, 2011.
- [24] Gandalf Lechner, Jan Schlemmer, and Yoh Tanimoto. On the equivalence of two deformation schemes in quantum field theory. *Lett. Math. Phys.*, to appear.
- [25] Gandalf Lechner and Christian Schützenhofer. Towards an operator-algebraic construction of integrable global gauge theories. *arXiv:1208.2366*, 2012.
- [26] Roberto Longo. Real Hilbert subspaces, modular theory,  $SL(2, \mathbf{R})$  and CFT. In *Von Neumann algebras in Sibiu: Conference Proceedings*, pages 33–91. Theta, Bucharest, 2008.
- [27] Roberto Longo and Edward Witten. An algebraic construction of boundary quantum field theory. *Comm. Math. Phys.*, 303(1):213–232, 2011.
- [28] S. N. M. Ruijsenaars. Scattering theory for the Federbush, massless Thirring and continuum Ising models. *J. Funct. Anal.*, 48(2):135–171, 1982.
- [29] S. N. M. Ruijsenaars. The Wightman axioms for the fermionic Federbush model. *Comm. Math. Phys.*, 87(2):181–228, 1982/83.
- [30] Bert Schroer. Localization and nonperturbative local quantum physics. *arXiv:hep-th/9805093*, 2012.
- [31] F. A. Smirnov. *Form factors in completely integrable models of quantum field theory*, volume 14 of *Advanced Series in Mathematical Physics*. World Scientific Publishing Co. Inc., River Edge, NJ, 1992.
- [32] M. Takesaki. *Theory of operator algebras. II*, volume 125 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2003. Operator Algebras and Non-commutative Geometry, 6.
- [33] Yoh Tanimoto. Construction of Wedge-Local Nets of Observables Through Longo-Witten Endomorphisms. *Comm. Math. Phys.*, 314(2):443–469, 2012.
- [34] Yoh Tanimoto. Noninteraction of Waves in Two-dimensional Conformal Field Theory. *Comm. Math. Phys.*, 314(2):419–441, 2012.
- [35] Mihály Weiner. An algebraic version of Haag’s theorem. *Comm. Math. Phys.*, 305(2):469–485, 2011.